arXiv:1504.05069v2 [math.QA] 29 Apr 2015

SUPER q-HOWE DUALITY AND WEB CATEGORIES

DANIEL TUBBENHAUER, PEDRO VAZ, AND PAUL WEDRICH

Abstract. We use super q-Howe duality to provide diagrammatic presentations of an idempotented form of the Hecke algebra and of categories of $\mathfrak{gl}_N$-modules (and more generally $\mathfrak{gl}_{N|M}$-modules) whose objects are tensor generated by exterior and symmetric powers of the vector representations. As an application, we give a representation theoretic explanation and a diagrammatic version of a symmetry of colored HOMFLY-PT polynomials.

Contents

1. Introduction 1
1.1. The framework 3
1.2. Outline of the paper 5
2. The diagrammatic categories 6
2.1. Definition of the category ∞-$\mathbf{Web}_{\mathrm{gr}}$ and its subquotients 6
2.2. The diagrammatic super relations 9
2.3. Green and red clasps 12
2.4. Braidings 14
2.5. A collection of diagrammatic idempotents 16
3. Proofs of the diagrammatic presentations 19
3.1. Super q-Howe duality 19
3.2. The sorted equivalences 23
3.3. Proofs of the equivalences 26
4. Applications 29
4.1. The colored HOMFLY-PT polynomial via the category ∞-$\mathbf{Web}_{\mathrm{gr}}$ 29
4.2. The colored $\mathfrak{sl}_N$-link polynomials via the categories N-$\mathbf{Web}_{\mathrm{gr}}$ 31
5. Generalization to webs for $\mathfrak{gl}_{N|M}$ 32
References 36

1. Introduction

Let $\mathbf{U}_q(\mathfrak{gl}_N)$ be the quantum enveloping $\mathbb{C}_q = \mathbb{C}(q)$-algebra for $\mathfrak{gl}_N$ with q being generic. Let $\mathfrak{gl}_N$-$\mathbf{Mod}_{\mathrm{es}}$ denote the braided monoidal category of $\mathbf{U}_q(\mathfrak{gl}_N)$-modules[1] tensor generated by *exterior* $\textstyle\bigwedge_q^k\mathbb{C}_q^N$ *and symmetric* $\mathrm{Sym}_q^l\mathbb{C}_q^N$ powers and $\mathbf{U}_q(\mathfrak{gl}_N)$-intertwiners between them.

We denote by $\check{\mathbf{H}}$ an *idempotented version* of the full collection of Iwahori-Hecke algebras $H_\infty(q) = \bigoplus_{K\in\mathbb{Z}_{\geq 0}} H_K(q)$ of type A. Roughly, $\check{\mathbf{H}}$ is the category obtained from the one-object

---

D.T. was supported by a research funding of the "Deutsche Forschungsgemeinschaft (DFG)" during the main part of this work. D.T. and P.W. thank the center of excellence grant "Centre for Quantum Geometry of Moduli Spaces (QGM)" from the "Danish National Research Foundation (DNRF)" for sponsoring a research visit which started this project. P.V. was financially supported by the "Université catholique de Louvain, Fonds Spéciaux de Recherche (FSR) 12 J.A.". P.W. was supported by an EPSRC doctoral training grant.

[1] We only consider finite dimensional, left modules throughout the paper.

category $H_{\infty}(q)$ by adding formal Gyoja-Aiston idempotents corresponding to column and row Young diagrams as new objects. By *quantum Schur-Weyl duality*, the categories $\mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}$ are quotients of $\check{\mathbf{H}}$ and the added idempotents can be thought of as lifts of the exterior $\bigwedge_q^k\mathbb{C}_q^N$ and the symmetric $\mathrm{Sym}_q^l\mathbb{C}_q^N$ powers.

We construct diagrammatic presentations of $\check{\mathbf{H}}$ and $\mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}$ by using the *green-red* web categories $\infty\text{-}\mathbf{Web}_{\mathrm{gr}}$ and $N\text{-}\mathbf{Web}_{\mathrm{gr}}$. Morphisms in these $\mathbb{C}_q$-linear categories are combinations of planar, upwards oriented, trivalent graphs with edges labeled by positive integers and colored black, green or red[2] modulo local relations. Objects are boundary conditions of such green-red webs, i.e. finite sequences of positive integers, each of which additionally carries a color black, green or red, indicated either by an actual coloring or by a subscript.

An example of a green-red web is:



A green integer k in a boundary sequence is meant to correspond to the $\mathbf{U}_q(\mathfrak{gl}_N)$-module $\bigwedge_q^k\mathbb{C}_q^N$, a red integer l to $\mathrm{Sym}_q^l\mathbb{C}_q^N$, and sequences of integers correspond to tensor products of such. Vertical edges are identities on these $\mathbf{U}_q(\mathfrak{gl}_N)$-modules and trivalent vertices encode more interesting $\mathbf{U}_q(\mathfrak{gl}_N)$-intertwiners. The integer 1 should be $\mathbb{C}_q^N$ independent of the color green or red, so we color it black.

Our main result is:

Theorem. (**The diagrammatic presentation**) The additive closures of $\infty\text{-}\mathbf{Web}_{\mathrm{gr}}$ and of $N\text{-}\mathbf{Web}_{\mathrm{gr}}$ are braided monoidally equivalent to $\check{\mathbf{H}}$ and $\mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}$ respectively.

We will see that $\infty\text{-}\mathbf{Web}_{\mathrm{gr}}$ admits an involution interchanging the colors green and red. An almost direct consequence of this is a symmetry between the HOMFLY-PT polynomials $\mathcal{P}^{a,q}(\cdot)$ of a link $\mathcal{L}$ whose strands are colored with Young diagrams $\vec{\lambda}=(\lambda^1,\dots,\lambda^d)$ or their transpose diagrams $\vec{\lambda}^{\,\mathrm{T}}=((\lambda^1)^{\mathrm{T}},\dots,(\lambda^d)^{\mathrm{T}})$.

Proposition. (**The colored HOMFLY-PT symmetry**) We have

$$\mathcal{P}^{a,q}(\mathcal{L}(\vec{\lambda}))\;=\;(-1)^{co}\,\mathcal{P}^{a,q^{-1}}(\mathcal{L}(\vec{\lambda}^{\,\mathrm{T}})). \tag{1}$$

Here *co* is some constant which only depends on the framed, oriented link $\mathcal{L}$ and its coloring.

Our results might help to understand symmetries observed within the homologies that categorify the colored HOMFLY-PT polynomials, see [10, Section 5].

Moreover, we show that a straightforward generalization of our approach also leads to diagrammatic presentations for categories $\mathfrak{gl}_{N|M}\text{-}\mathbf{Mod}_{\mathrm{es}}$ of $\mathbf{U}_q(\mathfrak{gl}_{N|M})$-modules tensor generated by exterior and symmetric powers of the vector representation. The presentations are given by quotients $N|M$-$\mathbf{Web}_{\mathrm{gr}}$ of ∞-$\mathbf{Web}_{\mathrm{gr}}$, which are obtained by killing Gyoja-Aiston idempotents corresponding to box-shaped Young diagrams.

---

[2] We use colored diagrams in this paper. The colors (black, green and red) are important and we recommend to read the paper in color. If the reader has a black-and-white version, then green will appear lightly shaded and black and red can be distinguished since black edges are always labeled 1.

1.1. **The framework.** One classical diagrammatic presentation is, modernly interpreted, due to Rumer, Teller and Weyl [24]. It states that the *Temperley-Lieb category* gives a diagrammatic presentation of the full subcategory of $\mathbf{U}_q(\mathfrak{sl}_2)$-modules tensor generated by the vector representation $\mathbb{C}_q^2$ of $\mathbf{U}_q(\mathfrak{sl}_2)$. In our notation, Temperley-Lieb diagrams can be seen as certain black-green webs with green edges of label 2 erased. Erasing of such edges is possible since they correspond to ${\bigwedge}_q^2\mathbb{C}_q^2 \cong \mathbb{C}_q$, which is the tensor unit in this representation category. The caps and cups in the Temperley-Lieb calculus can then be obtained because $\mathbb{C}_q^2 \cong (\mathbb{C}_q^2)^*$ holds as $\mathbf{U}_q(\mathfrak{sl}_2)$-modules.

Kuperberg extended this in [15] to all rank 2 Lie algebras. In particular, he gave a diagrammatic presentation[3] of the full subcategory of $\mathbf{U}_q(\mathfrak{sl}_3)$-modules tensor generated by the two *exterior powers* ${\bigwedge}_q^1\mathbb{C}_q^3 \cong \mathbb{C}_q^3$ and ${\bigwedge}_q^2\mathbb{C}_q^3$. His presentation can be recovered from ours by restricting to black-green webs, specializing $N = 3$ and the identification ${\bigwedge}_q^k\mathbb{C}_q^3 \cong ({\bigwedge}_q^{3-k}\mathbb{C}_q^3)^*$ of $\mathbf{U}_q(\mathfrak{sl}_3)$-modules.

Cautis, Kamnitzer and Morrison showed in [4] how to extend Kuperberg's ideas to give a diagrammatic presentation of $\mathfrak{gl}_N$-$\mathbf{Mod}_e$, the full subcategory of $\mathbf{U}_q(\mathfrak{gl}_N)$-modules tensor generated by the *exterior powers* ${\bigwedge}_q^k\mathbb{C}_q^N$ for $k = 0, \dots, N$. Partial results, specifically with respect to link polynomials, had been obtained before (for example in [20]). But the question remained open for a long time, since it was not clear how to prove completeness for a list of local relations in such a presentation.

The problem was solved in [4] by using *skew quantum Howe duality* (or short, *skew q-Howe duality*). In order to explain their approach, let $\vec{k} \in \mathbb{Z}_{\geq 0}^m$ such that $k_1 + \dots + k_m = K$. By skew q-Howe duality, the commuting actions of $\mathbf{U}_q(\mathfrak{gl}_m)$ and $\mathbf{U}_q(\mathfrak{gl}_N)$ on

$$ {\bigwedge}_q^K(\mathbb{C}_q^m \otimes \mathbb{C}_q^N) \cong \textstyle\bigoplus_{\vec{k}\in\mathbb{Z}_{\geq 0}^m} {\bigwedge}_q^{k_1}\mathbb{C}_q^N \otimes \dots \otimes {\bigwedge}_q^{k_m}\mathbb{C}_q^N $$

give rise to a functor $\Phi_{\mathrm{skew}}^m \colon \dot{\mathbf{U}}_q(\mathfrak{gl}_m) \to \mathfrak{gl}_N$-$\mathbf{Mod}_e$, where $\dot{\mathbf{U}}_q(\mathfrak{gl}_m)$ is the idempotented form of $\mathbf{U}_q(\mathfrak{gl}_m)$. Then Cautis, Kamnitzer and Morrison considered, in our notation[4],

$$\begin{tikzcd}
\dot{\mathbf{U}}_q(\mathfrak{gl}_m) \arrow[r, "\Phi_{\mathrm{skew}}^m"] \arrow[dr, "\Upsilon_{\mathrm{skew}}^m"'] & \mathfrak{gl}_N\text{-}\mathbf{Mod}_e \\
 & N\text{-}\mathbf{Web}_{\mathrm{g}} \arrow[u, "\Gamma"']
\end{tikzcd} , \tag{2}$$

where $\Upsilon_{\mathrm{skew}}^m$ is a certain *ladder functor* realizing an action of $\dot{\mathbf{U}}_q(\mathfrak{gl}_m)$ on the diagram category N-$\mathbf{Web}_{\mathrm{g}}$. The *presentation functor* Γ is constructed such that (2) commutes. The functor Φ_{skew}^m is full and it remains to study its kernel, which, in this case, is only generated by killing $\mathfrak{gl}_m$-weights with entries not in $\{0, \dots, N\}$. That Γ is an equivalence follows since N-$\mathbf{Web}_{\mathrm{g}}$ is defined to be the quotient of a "free" web category by relations coming from $\dot{\mathbf{U}}_q(\mathfrak{gl}_m)$ (to make the ladder functor $\Upsilon^m_{\mathrm{skew}}$ well-defined) and the $\Upsilon^m_{\mathrm{skew}}$ image of the kernel of Φ^m_{skew}. As before, $\mathfrak{sl}_N$-$\mathbf{Mod}_e$ can be recovered by identifying $\textstyle\bigwedge^k_q\mathbb{C}^N_q \cong (\textstyle\bigwedge^{N-k}_q\mathbb{C}^N_q)^*$ as $\mathbf{U}_q(\mathfrak{sl}_N)$-modules.

---

[3] In fact, Kuperberg as well as [4] worked more generally with spiders and pivotal equivalences. Moreover, also [23] considered a more general framework by allowing caps and cups.

[4] We consider $\mathfrak{gl}_N$-$\mathbf{Mod}_{\mathrm{es}}$ instead of $\mathfrak{sl}_N$-$\mathbf{Mod}_{\mathrm{es}}$, see also Remark 1.1.

In [23] the situation of *symmetric quantum Howe duality* (for short, symmetric q-Howe duality) was studied[5]. That is, there is an analogue of (2) where $\mathfrak{gl}_N$-$\mathbf{Mod}_e$ is replaced by $\mathfrak{gl}_N$-$\mathbf{Mod}_s$, the full subcategory $\mathbf{U}_q(\mathfrak{gl}_N)$-modules tensor generated by the *symmetric powers* $\mathrm{Sym}^l_q\mathbb{C}^N_q$ for $l \in \mathbb{Z}_{\geq 0}$. Moreover, N-$\mathbf{Web}_{\mathrm{g}}$ is replaced by N-$\mathbf{Web}_{\mathrm{r}}$ which can be recovered from our diagrammatic presentation by considering purely red webs. Again, it remained to study the kernel of the corresponding functor Φ^m_{sym} and this was successfully done in [23] for $N = 2$. In this case, the kernel of Φ^m_{sym} is generated by killing $\mathfrak{gl}_m$-weights with negative entries and one additional *dumbbell relation*, which encodes the $\mathfrak{gl}_2$-$\mathbf{Mod}_s$ relation $\mathbb{C}^2_q \otimes \mathbb{C}^2_q \cong \mathbb{C}_q \oplus \mathrm{Sym}^2_q\mathbb{C}^2_q$. A direct generalization for $N > 2$ would require even more extra relations, besides killing $\mathfrak{gl}_m$-weights.

In this paper we give a diagrammatic presentation of the category $\mathfrak{gl}_N$-$\mathbf{Mod}_{\mathrm{es}}$, a category of $\mathbf{U}_q(\mathfrak{gl}_N)$-modules containing both exterior *and* symmetric powers of the vector representation. This diagrammatic presentation gives a common generalization of the web categories of [4] and [23]. Our approach is to *turn Cautis, Kamnitzer and Morrison's recipe on the head*: we see their approach as a *machine* that "takes dualities and produces diagrammatic presentations of the related representation theoretical categories". Specifically, we start with *super quantum Howe duality* (for short, super q-Howe duality) between the superalgebra $\mathbf{U}_q(\mathfrak{gl}_{m|n})$ and $\mathbf{U}_q(\mathfrak{gl}_N)$. We obtain a full super q-Howe functor $\Phi^{m|n}_{\mathrm{su}}$, which we attempt to factor as a composite of a ladder functor $\Upsilon^{m|n}_{\mathrm{su}}$ – mapping into an appropriate web category – and a diagrammatic presentation functor Γ_N, to give an analogue of the commutative diagram (2). In this analogue $\dot{\mathbf{U}}_q(\mathfrak{gl}_m)$ is replaced by $\dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n})$, $\mathfrak{gl}_N$-$\mathbf{Mod}_e$ is replaced by a category of $\mathbf{U}_q(\mathfrak{gl}_N)$-modules tensor generated by exterior $\textstyle\bigwedge^k_q\mathbb{C}^N_q$ and symmetric powers $\mathrm{Sym}^l_q\mathbb{C}^N_q$, and N-$\mathbf{Web}_{\mathrm{g}}$ is replaced by a category of green-red webs, where the color indicates whether an edge represents an exterior or a symmetric power. Having decided to follow this strategy, the definition of the appropriate web category – as a quotient of a "free" green-red web category – is already determined. Two aspects are important:

(I) In order to make $\Upsilon^{m|n}_{\mathrm{su}}$ well-defined, the web category needs to satisfy ladder images of $\dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n})$ relations. Remarkably, it suffices to consider relations coming from the subalgebra $\dot{\mathbf{U}}_q(\mathfrak{gl}_m) \oplus \dot{\mathbf{U}}_q(\mathfrak{gl}_n)$ and only one additional *super commutation* relation $[2]1_{\vec{k}} = F_m E_m 1_{\vec{k}} + E_m F_m 1_{\vec{k}}$ for $\mathfrak{gl}_{m|n}$-weights with $k_m = k_{m+1} = 1$. This corresponds to the *dumbbell relation* on webs and to $\mathbb{C}^N_q \otimes \mathbb{C}^N_q \cong \textstyle\bigwedge^2_q\mathbb{C}^N_q \oplus \mathrm{Sym}^2_q\mathbb{C}^N_q$ in $\mathfrak{gl}_N$-$\mathbf{Mod}_{\mathrm{es}}$.

(II) In order to make the diagrammatic presentation functor an equivalence, we need to impose the ladder image of $\ker(\Phi^{m|n}_{\mathrm{su}})$ as relations in the web category. In fact, $\ker(\Phi^{m|n}_{\mathrm{su}})$ is spanned by idempotents corresponding to $\mathfrak{gl}_{m|n}$-weights $\vec{k} = (k_1, \dots, k_{m+n})$ with $k_1, \dots, k_m \notin \{0, \dots, N\}$ or $k_{m+1}, \dots, k_{m+n} \notin \mathbb{Z}_{\geq 0}$. It is remarkable that no extra relations, aside from killing these $\mathfrak{gl}_{m|n}$-weights, are necessary.

---

[5] In fact, the observations made in the paper [23] were one of the main motivations to start this project and are used throughout the present paper in many ways.

We impose the ladder images of $\ker(\Phi^{m|n}_{\mathrm{su}})$ in two steps: first we kill all $\mathfrak{gl}_{m|n}$-weights with negative entries by allowing only non-negative labels on web edges. This produces the web category ∞-$\mathbf{Web}_{\mathrm{gr}}$, which is symmetric under exchanging green and red. On this we further quotient by setting $\mathfrak{gl}_{m|n}$-weights $\vec{k} = (k_1, \dots, k_{m+n})$ to zero if one of $k_1, \dots, k_m$ is greater than N. This produces the web category N-$\mathbf{Web}_{\mathrm{gr}}$ and in Theorem 3.20 we show that its additive closure is equivalent to $\mathfrak{gl}_N$-$\mathbf{Mod}_{\mathrm{es}}$. In Theorem 3.22 we use *quantum Schur-Weyl duality* to derive from Theorem 3.20 that ∞-$\mathbf{Web}_{\mathrm{gr}}$ gives a diagrammatic presentation of the idempotented Hecke algebra $\check{\mathbf{H}}$ from above.

Remark 1.1. We describe $\mathfrak{gl}_N$-$\mathbf{Mod}_{\mathrm{es}}$ and not $\mathfrak{sl}_N$-$\mathbf{Mod}_{\mathrm{es}}$ because of the algebraic form of super q-Howe duality. In particular, our web categories do not contain duality isomorphisms $\bigwedge^k_q \mathbb{C}^N_q \cong (\bigwedge^{N-k}_q \mathbb{C}^N_q)^*$, which would be necessary for a diagrammatic presentation of $\mathfrak{sl}_N$-$\mathbf{Mod}_{\mathrm{es}}$. In $\mathfrak{gl}_N$-$\mathbf{Mod}_{\mathrm{es}}$, on the other hand, there are no such hidden duals, as we have $\bigwedge^k_q \mathbb{C}^N_q \cong \bigwedge^N_q \mathbb{C}^N_q \otimes (\bigwedge^{N-k}_q \mathbb{C}^N_q)^*$ as $\mathbf{U}_q(\mathfrak{gl}_N)$-modules. Here $\bigwedge^N_q \mathbb{C}^N_q \cong L((1, \dots, 1))$ is the $\mathbf{U}_q(\mathfrak{gl}_N)$-module of highest weight $\lambda = (1, \dots, 1) \in \mathbb{Z}^N_{\geq 0}$.

Last, but not least, we use the more general *super q-Howe duality between* $\mathbf{U}_q(\mathfrak{gl}_{m|n})$ *and* $\mathbf{U}_q(\mathfrak{gl}_{N|M})$ to describe $\mathfrak{gl}_{N|M}$-$\mathbf{Mod}_{\mathrm{es}}$. Feeding that duality into the "diagrammatic presentation machine" shows that this representation category is equivalent to the quotient $N|M$-$\mathbf{Web}_{\mathrm{gr}}$ of ∞-$\mathbf{Web}_{\mathrm{gr}}$, which is obtained by killing the Gyoja-Aiston idempotent corresponding to the size $(N+1) \times (M+1)$ box-shaped Young diagram. This is a generalization, since for $M = 0$, $\mathfrak{gl}_{N|M}$-$\mathbf{Mod}_{\mathrm{es}}$ is equivalent to $\mathfrak{gl}_N$-$\mathbf{Mod}_{\mathrm{es}}$ and $N|M$-$\mathbf{Web}_{\mathrm{gr}}$ is equal to N-$\mathbf{Web}_{\mathrm{gr}}$, because the box idempotent corresponds exactly to an $(N+1)$-labeled green edge.

This diagrammatic presentation generalizes Grant's and Sartori's presentations of the category $\mathfrak{gl}_{1|1}$-$\mathbf{Mod}_{\mathrm{e}}$, see [8] and [25], and the diagrammatic calculus for $\mathfrak{gl}_{N|M}$-$\mathbf{Mod}_{\mathrm{e}}$ given in [21] (see also [7]). Compared to the latter, our generalization, which also takes the symmetric powers of $\mathbb{C}^{N|M}_q$ into account, does not need any extra relations aside from the dumbbell relation. In fact, the one extra relation needed to make the diagrammatic calculus given in [21] faithful, see [21, Remark 6.19], has a very compact and natural description in our green-red web category $N|M$-$\mathbf{Web}_{\mathrm{gr}}$.

Finally, we sketch how our diagrammatic presentation of $\mathfrak{gl}_{N|M}$-$\mathbf{Mod}_{\mathrm{es}}$ extends to take duals of exterior and symmetric powers into account. This closely follows [21, Section 6]. The resulting diagrammatic category allows the computation of the colored Reshetikhin-Turaev $\mathfrak{gl}_{N|M}$-link invariants. In Corollary 5.13, we interpret the colored HOMFLY-PT symmetry (1) as a stable version of a symmetry between colored Reshetikhin-Turaev $\mathfrak{gl}_{N|M}$- and $\mathfrak{gl}_{M|N}$-link invariants.

1.2. **Outline of the paper.** Section 2 is the diagrammatic heart of our paper where we introduce ∞-$\mathbf{Web}_{\mathrm{gr}}$ and it subquotients N-$\mathbf{Web}_{\mathrm{gr}}$, N-$\mathbf{Web}_{\mathrm{g}}$ and N-$\mathbf{Web}_{\mathrm{r}}$.

Section 3 contains the proof of our main theorems and splits into three subsections: we first introduce super q-Howe duality. Then we show an equivalence between "sorted" subcategories of N-$\mathbf{Web}_{\mathrm{gr}}$ and $\mathfrak{gl}_N$-$\mathbf{Mod}_{\mathrm{es}}$. These subcategories are induced by the algebraic form of super q-Howe duality. By using the "sorted" equivalence and the fact that the braiding gives a way to "shuffle" the "sorted" subcategories, we prove our main theorems.

In Section 4 we discuss one application of our diagrammatic presentation: we give a procedure to recover the colored HOMFLY-PT polynomial from ∞-$\mathbf{Web}_{\mathrm{gr}}$. A direct consequence of the green-red symmetry is a symmetry within the colored HOMFLY-PT polynomial obtained by transposing Young diagrams, see (1). The colored Reshetikhin-Turaev $\mathfrak{sl}_N$-link polynomials can be recovered from our approach as well, as we sketch in the last subsection.

Finally, in Section 5 we generalize the diagrammatic presentation of $\mathfrak{gl}_N$-$\mathbf{Mod}_{\mathrm{es}}$ to the super case $\mathfrak{gl}_{N|M}$-$\mathbf{Mod}_{\mathrm{es}}$ and we sketch an extension of our diagrammatic calculus to include dual representations. The required arguments are – mutatis mutandis – contained in the previous sections and in [21, Section 6], which allows a very compact exposition in Section 5.

Acknowledgements: We thank Antonio Sartori for a careful reading of a draft version of this paper and many helpful comments. Special thanks to David Rose – many of the ideas underlying this paper came up in the joint work between him and D.T. We also thank Jonathan Brundan, Jonathan Grant, Jonathan Kujawa, Marco Mackaay, Weiwei Pan, Jake Rasmussen, Marko Stošić, Catharina Stroppel, Geordie Williamson and Oded Yacobi for helpful discussion, comments, and probing questions. We also like to thank Skype for many useful conversations.

2. The diagrammatic categories

In the present section we introduce the category ∞-$\mathbf{Web}_{\mathrm{gr}}$ and its quotient N-$\mathbf{Web}_{\mathrm{gr}}$. These provide diagrammatic presentations of $\check{\mathbf{H}}$ and its quotient categories $\mathfrak{gl}_N$-$\mathbf{Mod}_{\mathrm{es}}$ respectively. Other subquotients of ∞-$\mathbf{Web}_{\mathrm{gr}}$ are N-$\mathbf{Web}_{\mathrm{g}}$ and N-$\mathbf{Web}_{\mathrm{r}}$ (and later in Section 5, $N|M$-$\mathbf{Web}_{\mathrm{gr}}$) which are related to categories studied in [4] and in [23] respectively.

2.1. Definition of the category ∞-$\mathbf{Web}_{\mathrm{gr}}$ and its subquotients.

We first introduce the *free green-red web category* ∞-$\mathbf{Web}^{f}_{\mathrm{gr}}$. To this end, we denote by X the set

$$X = X_b \cup X_g \cup X_r = \{0_b, 1_b\} \cup \{2_g, 3_g, \dots\} \cup \{2_r, 3_r, \dots\},$$

where we think of the elements of X_b as being colored *black*, of the elements of X_g as being colored *green* and of the elements of X_r as being colored *red*. We usually omit the subscripts, since the colors on the boundary can be read off from the diagrams.

Definition 2.1. The *free green-red web category*, which we denote by ∞-$\mathbf{Web}^{f}_{\mathrm{gr}}$, is the category determined by the following data.

- The objects of ∞-$\mathbf{Web}^{f}_{\mathrm{gr}}$ are finite (possibly empty) sequences $\vec{k} \in X^L$ with entries from X for some $L \in \mathbb{Z}_{\geq 0}$, together with a zero object. We display the entries of $\vec{k}$ ordered from left to right according to their appearance in $\vec{k}$.
- The morphism space $\mathrm{Hom}_{\infty\text{-}\mathbf{Web}^{f}_{\mathrm{gr}}}(\vec{k}, \vec{l})$ from $\vec{k}$ to $\vec{l}$ is the $\mathbb{C}_q$-vector space spanned by isotopy classes[6] of planar, upwards oriented, trivalent graphs with edges labeled by positive integers and colored black, green or red, with bottom boundary $\vec{k}$ and top boundary $\vec{l}$. More precisely, we only allow webs that can be obtained by composition $\circ$ (vertical gluing) and tensoring $\otimes$ (horizontal juxtaposition) of the following basic pieces (including the empty diagram).

---

[6] We require that isotopies preserve the upward orientations and the boundary of green-red webs.

Let $k, l \in \mathbb{Z}_{\geq 2}$, then the generators are



called (from left to right) *black identity*, *green identity*, *red identity*, *green merge*, *green split*, *red merge* and *red split*, together with (here $k, l \in \mathbb{Z}_{\geq 0}$)

(3)



called *mixed merges* and *mixed splits* respectively.

We call webs obtained by composition of generators with only black and green edges or only black and red edges *monochromatic*. ◇

Remark 2.2. Note the following conventions and properties of ∞-$\mathbf{Web}^{f}_{\mathrm{gr}}$.

- The category is $\mathbb{C}_q$-linear, i.e. the spaces $\mathrm{Hom}_{\infty\text{-}\mathbf{Web}^{f}_{\mathrm{gr}}}(\vec{k}, \vec{l})$ are $\mathbb{C}_q$-vector spaces and the composition $\circ$ is $\mathbb{C}_q$-bilinear. Moreover, the category is monoidal by juxtaposition $\otimes$ of objects and morphisms. $\otimes$ is also $\mathbb{C}_q$-bilinear on morphism spaces.
- As a shorthand notation, we allow edges of label 1 to have any color:

(4)



and



although formally green and red edges with label 1 do not exist. The merges and splits on the right are obtained by setting $k = 1$ or $l = 1$ in (3).
- The reading conventions for all webs are from *bottom to top* and *left to right*: given two webs u, v, then $v \circ u$ is obtained by gluing v on top of u and $u \otimes v$ is given by putting v to the right of u. Moreover, if any of the top boundary labels of u differs from the corresponding bottom boundary label of v, then, by convention, $v \circ u = 0$.
- For $j \in \mathbb{Z}_{\geq 1}$, we define the so-called *monochromatic* $F^{(j)}1_{(k,l)}$ *and* $E^{(j)}1_{(k,l)}$*-ladders* as

(5) $F^{(j)}1_{(k,l)} =$



, $E^{(j)}1_{(k,l)} =$



and analogously in red. Sometimes we draw such ladder rungs horizontally. We also have the *mixed* $F1_{(k,l)}$ *and* $E1_{(k,l)}$*-ladders*

(6) $F1_{(k,l)} =$



, $E1_{(k,l)} =$



Similarly by exchanging green and red. Note that the ladders from (5) exist for all $j \in \mathbb{Z}_{\geq 1}$, while the mixed ladders from (6) exist only for $j = 1$.

Definition 2.3. The *green-red web category* ∞-$\mathbf{Web}_{\mathrm{gr}}$ is the quotient of ∞-$\mathbf{Web}_{\mathrm{gr}}^{f}$ obtained by imposing the following local relations on morphisms. The *monochromatic relations*, which hold for green webs as well as for red webs: *associativity*

(7)



where we use the shorthand notation from (4) if some of the labels are 1. Next, the *digon removal relations*

(8)



for which k, l might be 1. In these relations the *s,t-quantum binomial* is given by

$$\begin{bmatrix} s \\ t \end{bmatrix} = \frac{[s][s-1]\cdots[s-t+2][s-t+1]}{[t]!} \in \mathbb{C}_q.$$

Here $[s] = \frac{q^s - q^{-s}}{q - q^{-1}} \in \mathbb{C}_q$ is the *s-th quantum number* and $[t]! = [1][2]\cdots[t-1][t] \in \mathbb{C}_q$ is the *t-th quantum factorial* ($s \in \mathbb{Z}$, $t \in \mathbb{Z}_{\geq 0}$). Finally, the (in)famous *square switch* relations

(9)



Here we allow j_1 or j_2 to be 1 (we will get *mixed* square switch relations, with one green and one red side, in Lemma 2.10).

The defining relation between green and red edges is the *dumbbell relation*:

(10)



In addition to the above: webs with edges labeled < 0 do not exist in our calculus. Since it is sometimes convenient in pictures to illustrate webs that could potentially have edges labeled < 0, we define these to be zero. ◇

Remark 2.4. Observe that ∞-$\mathbf{Web}_{\mathrm{gr}}$ is symmetric under exchanging green and red. In the following we will often refer to this symmetry to shorten arguments.

Definition 2.5. The category N-$\mathbf{Web}_{\mathrm{gr}}$ is the quotient category obtained from ∞-$\mathbf{Web}_{\mathrm{gr}}$ by imposing the *exterior relations*, that is,

(11) 

$$= 0 \quad , \quad \text{if } k > N.$$

The exterior relations hold only for green edges. These relations mean that any web u with a green edge labeled $k > N$ is zero. In contrast, red edges labeled $k > N$ are usually not zero.

The sorted web category N-$\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}}$ is the full (non-monoidal) subcategory of N-$\mathbf{Web}_{\mathrm{gr}}$ whose object set consists of $\vec{k} \in X^L$ with no red boundary point left of a green boundary point: if $k_i \in X_r$ for some i, then $k_{>i} \in X_b \cup X_r$. ◇

Remark 2.6. The relations (11) are diagrammatic versions of

$$\textstyle\bigwedge_q^{>N} \mathbb{C}_q^N \cong 0.$$

Definition 2.7. The category N-$\mathbf{Web}_{\mathrm{g}}$ is the subcategory of N-$\mathbf{Web}_{\mathrm{gr}}$ consisting of only black and green objects and whose morphism spaces are spanned as $\mathbb{C}_q$-vector spaces by webs that contain only black or green edges.

Similarly, the category N-$\mathbf{Web}_{\mathrm{r}}$ is the subcategory of N-$\mathbf{Web}_{\mathrm{gr}}$ consisting of only black and red objects and whose morphism spaces are spanned as $\mathbb{C}_q$-vector spaces by webs that contain only black or red edges.

We call these categories *monochromatic.* ◇

Remark 2.8. We will see in Corollary 2.16 that N-$\mathbf{Web}_{\mathrm{g}}$ is equivalent to the web category given in [4, Definition 2.2] (without tags and downwards pointing arrows). The category N-$\mathbf{Web}_{\mathrm{r}}$ is a generalization of the one given in [23, Definition 1.4]. In fact, Proposition 2.15 shows that both monochromatic subcategories are full in N-$\mathbf{Web}_{\mathrm{gr}}$.

2.2. **The diagrammatic super relations.** We show in this subsection that diagrammatic versions of the relations (16) in the Howe dual quantum group $\dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n})$ from Definition 3.1 hold in our diagrammatic categories ∞-$\mathbf{Web}_{\mathrm{gr}}$ and N-$\mathbf{Web}_{\mathrm{gr}}$.

Lemma 2.9. We have the relations



$$= 0 =$$


where the dots should indicate k parallel black edges with label 1 which split off the bottom and merge with the top in any order (the order does not matter because of (7)).

Proof. It suffices by associativity (7) to show the statement for $k = 2$. We have



The other $k = 2$ relation follows by symmetry. ☐

Lemma 2.10. (a) We have for all $k, l \in \mathbb{Z}_{\geq 0}$



(b) We have for all $k, l \in \mathbb{Z}_{\geq 0}$



Similarly for exchanged roles of green and red.

(c) We have for all $k, l \in \mathbb{Z}_{\geq 0}$



Similarly for exchanged roles of green and red and flipped horizontal orientations.

Proof. **(a)**: Directly from (7), Lemma 2.9 and symmetry.

(b): Let u and v denote the two webs on the right-hand side of (b) above. Using (9) for the edges labeled $k+1$ and $l+1$ in u respectively v, we get



after collapsing appearing digons. By using (10) on the central vertical edges in the expansions, we see that $u + v = s \cdot \mathrm{id}_{(k,l)}$. The scalar is $s = [2][k][l] + [k][1-l] - [k-1][l] = [k+l]$. The other cases follow by symmetry.

(c): We start with the web on the left-hand side and first use (10) on the middle two horizontal edges. Thus, we obtain (our drawings are simplified and the orientations pointing down could be isotoped to point up)



The two marked parts above are monochromatic squares, which can be switched to give:



Plugging these four terms back in, we get the four webs from the right-hand side of the equation in (c) (in the indicated order) which can be seen by using (7) as for example



The other three cases in (c) follow by symmetry. ☐

2.3. Green and red clasps.

We show now that our calculus contains web analogues of the *Jones-Wenzl projectors* of the Temperley-Lieb algebra. We call them *clasps*, following [15].

From now on, we denote by capital vectors as $\vec{K} \in X^K$ special objects of ∞-**Web**$_{\mathrm{gr}}$ of the form $\vec{K} = (1_b, \dots, 1_b)$ with K entries equal 1_b and no other entries.

Definition 2.11. Let $K \in \mathbb{Z}_{>0}$. We define the *K-th green clasp* $\mathcal{CL}^g_K \in \mathrm{End}_{\infty\text{-}\mathbf{Web}_{\mathrm{gr}}}(\vec{K})$ recursively: $\mathcal{CL}^g_1$ is the black identity strand and for $K \in \mathbb{Z}_{>1}$ set



Similarly for the *red clasp* $\mathcal{CL}^r_K$ by exchanging green and red. ◇

The following lemma identifies the clasps avoiding the recursive definition.

Lemma 2.12. We have for all $K \in \mathbb{Z}_{>0}$


$\mathcal{CL}^g_K = \dfrac{1}{[K]!}$
1 · · · 1
K
1 · · · 1
,
$\mathcal{CL}^r_K = \dfrac{1}{[K]!}$
1 · · · 1
K
1 · · · 1

where we repeatedly split an edge labeled K until all of the top and bottom edges are black.

Proof. Up to signs and drawing conventions as in [23, Lemma 2.12] and left to the reader. ☐

Corollary 2.13. For all $K \in \mathbb{Z}_{>0}$: the projector $\mathcal{CL}_K^g$ can be expressed as a linear combination of webs with only black and red edges of label 2. Similarly for $\mathcal{CL}_K^r$.

Proof. Directly from (10) and Lemma 2.12. □

Example 2.14. The projector $\mathcal{CL}_1^r$ is just the black identity strand, the projector $\mathcal{CL}_2^r$ is $\frac{1}{[2]}$ times the red dumbbell as in (10) and



Note that all edges appearing on the right-hand side are black or green with label 2. ■

Proposition 2.15. Let $\vec{k}$ and $\vec{l}$ be sequences of black and green boundary points. Every web $u \in \mathrm{Hom}_{\infty\text{-}\mathbf{Web}_{\mathrm{gr}}}(\vec{k}, \vec{l})$ can be expressed as a sum of webs with only black and green edges. Similarly by exchanging green and red.

Proof. We start by exploding[7] every red edge. Around internal vertices of u with no outgoing green edges we get



Note that the marked part above is $\mathcal{CL}_{k+l}^r$ up to a non-zero scalar. Thus, we can use Corollary 2.13 to replace $\mathcal{CL}_{k+l}^r$ by a non-zero sum of webs with only black and green edges. Repeating this for all purely red internal vertices shows the statement, since all outer edges are assumed to be black or green. The other statement follows by symmetry. □

Denote by N-$\mathbf{Web}_{\mathrm{CKM}}$ the category given in [4, Definition 2.2] (without tags and downwards pointing arrows). As a consequence of Proposition 2.15 we see that interpreting webs in N-$\mathbf{Web}_{\mathrm{CKM}}$ as green webs in N-$\mathbf{Web}_{\mathrm{gr}}$ gives a full functor ι_1^∞ between these categories. In Lemma 3.13 we will see that it is also faithful and we get the following corollary.

Corollary 2.16. The functor $\iota_1^\infty \colon N\text{-}\mathbf{Web}_{\mathrm{CKM}} \to N\text{-}\mathbf{Web}_{\mathrm{gr}}$, given by coloring webs green, is an inclusion of a full, monoidal subcategory. In particular, N-$\mathbf{Web}_{\mathrm{CKM}}$ and N-$\mathbf{Web}_{\mathrm{g}}$ are equivalent as monoidal categories.

Proof. The functor is well-defined since all relations in N-$\mathbf{Web}_{\mathrm{CKM}}$ hold in N-$\mathbf{Web}_{\mathrm{gr}}$. That ι_1^∞ is monoidal is clear, fullness follows from Proposition 2.15 and faithfulness from Lemma 3.13. Thus, we see that N-$\mathbf{Web}_{\mathrm{CKM}}$ and N-$\mathbf{Web}_{\mathrm{g}}$ are monoidally equivalent. □

---

[7] We "explode" by using (8) (the order of does not matter by (7)). We indicate "explosions" with dots.

2.4. **Braidings.** We define now a *braided* monoidal structure on ∞-$\mathbf{Web}_{\mathrm{gr}}$.

Definition 2.17. Define for $k,l\in\mathbb{Z}_{\geq 0}$ an *elementary crossing* depending on four cases. The *monochromatic crossings* (note the different powers of q):

(12)

$\beta^{g}_{k,l} =$ 

$= (-1)^{k+kl}q^{k} \sum_{\substack{j_1,j_2\geq 0\\ j_1-j_2=k-l}} (-q)^{-j_1}$ 

$\beta^{r}_{k,l} =$ 

$= (-1)^{k}q^{-k} \sum_{\substack{j_1,j_2\geq 0\\ j_1-j_2=k-l}} (-q)^{+j_1}$ 

The *mixed crossings* are defined via explosion of the strand going over:

(13) $\beta^{m}_{k,l} =$ 

$= \frac{1}{[k]!}$ 

and $\beta^{\tilde{m}}_{k,l} =$ 

$= \frac{1}{[k]!}$ 

where the remaining crossings are of the form $\beta^{r}_{1,l}$ or $\beta^{g}_{1,l}$ respectively. ◇

Example 2.18. The case $k=l=1$ is not ambiguous, since we have

$\beta^{g}_{1,1} = q\Bigg($ 

$- q^{-1}$ 

$\Bigg) \overset{(10)}{=} -q^{-1}\Bigg($ 

$- q$ 

$\Bigg) = \beta^{r}_{1,1},$

as a small calculation shows. ■

As a shorthand notation, we write $\beta^{\bullet}_{k,l}$ where $\bullet$ stands for either g, r, m or $\tilde{m}$ from now on. Note that the sums in (12) are finite, because webs with negative labels are zero.

Lemma 2.19. (**Pitchfork relations**) We have



$=$ 

, 

$=$ 

Similar with exchanged roles of green and red, for the monochromatic cases and with merges.

Note that the pitchfork lemma directly implies that (13) could also be done by exploding the edges going underneath instead of the edges going over (or exploding both).

Proof. The pitchfork lemma with only green colored edges follows as in [20, Lemma 5.3]. By symmetry, the arguments go through for the monochromatic red case as well.

The mixed, left-hand equation is easy to verify by the above, since we explode the over-crossing edge and we thus, can directly use the monochromatic case. It remains to prove the mixed, right-hand equation. We only need to check the case $k = 2$, the case $k \in \mathbb{Z}_{>2}$ then follows easily from this case by using Lemma 2.9. We write



The rightmost diagram is zero by Lemma 2.9 and the monochromatic pitchfork relations. This proves the mixed right-hand equation. The other cases are analogous. ☐

Let $\vec{k} \in X^{L}_{\geq 0}$ be an object in ∞-**Web**$_{\mathrm{gr}}$. We define for $i = 1, \dots, L-1$ the crossing $\beta^{\bullet}_{i} 1_{\vec{k}}$ to be the corresponding elementary crossing $\beta^{\bullet}_{k_i,k_{i+1}}$ between the strands i and $i+1$ and the identity elsewhere. Clearly, it suffices to indicate the rightmost $1_{\vec{k}}$ in a sequence of $\beta^{\bullet}_{i} 1_{\vec{k}}$'s.

Lemma 2.20. The crossings $\beta^{\bullet}_{i} 1_{\vec{k}}$ satisfy the braid relations, that is, they are invertible, they satisfy the commutation relations $\beta^{\bullet}_{i}\beta^{\bullet}_{j} 1_{\vec{k}} = \beta^{\bullet}_{j}\beta^{\bullet}_{i} 1_{\vec{k}}$ for $|i-j| > 2$ and the Reidemeister 3 relations $\beta^{\bullet}_{i}\beta^{\bullet}_{j}\beta^{\bullet}_{i} 1_{\vec{k}} = \beta^{\bullet}_{j}\beta^{\bullet}_{i}\beta^{\bullet}_{j} 1_{\vec{k}}$ for $|i-j| = 1$.

The inverses $\beta^{\bullet -1}_{i}$ are given as in (12), but with $q \to q^{-1}$. See also [20, Section 5].

Proof. By Lemma 2.19, since the black case can be verified as in [20, Section 5]. ☐

Remark 2.21. Let S_K denote the symmetric group in K letters. Moreover, let $w \in S_K$ and let $\beta^{\bullet}_{w} \in \mathrm{End}_{\infty\text{-}\mathbf{Web}_{\mathrm{gr}}}(\vec{K})$ be the permutation braid associated to w (this is a well-defined assignment by Lemma 2.20). Let $\ell(w)$ be the length of w. Following [14, Chapter 3, Section 2], one can show that

$$\mathcal{CL}^{g}_{K} = q^{\frac{K(K-1)}{2}} \frac{1}{[K]!} \sum_{w \in S_K} (-q)^{-\ell(w)} \beta^{\bullet}_{w} \quad , \quad \mathcal{CL}^{r}_{K} = q^{-\frac{K(K-1)}{2}} \frac{1}{[K]!} \sum_{w \in S_K} q^{\ell(w)} \beta^{\bullet}_{w}.$$

The factors $q^{\frac{K(K-1)}{2}}$ and $q^{-\frac{K(K-1)}{2}}$ come from our conventions for crossings.

Define $\beta^{\bullet}_{\vec{k},\vec{l}}$ for objects $\vec{k} = (k_1, \dots, k_a)$ and $\vec{l} = (l_1, \dots, l_b)$ via

$\beta^{\bullet}_{\vec{k},\vec{l}} =$


l_1 ··· l_b k_1 ··· k_a
k_1 ··· k_a l_1 ··· l_b

$\in \mathrm{Hom}_{\infty\text{-}\mathbf{Web}_{\mathrm{gr}}}(\vec{k} \otimes \vec{l}, \vec{l} \otimes \vec{k}),$

where blue should stand for all suitable color possibilities.

Recall that a *braided monoidal category* (with an underlying strict monoidal category) is a pair $(\boldsymbol{\mathcal{C}}, \beta^{\boldsymbol{\mathcal{C}}}_{\cdot,\cdot})$ consisting of a monoidal category $\boldsymbol{\mathcal{C}}$ and a collection of natural isomorphisms $\beta^{\boldsymbol{\mathcal{C}}}_{\vec{k},\vec{l}} \colon \vec{k} \otimes \vec{l} \to \vec{l} \otimes \vec{k}$ such that the *hexagon identities* hold for any objects $\vec{k}, \vec{l}, \vec{m}$ of $\boldsymbol{\mathcal{C}}$:

$$\beta^{\boldsymbol{\mathcal{C}}}_{\vec{k},\vec{l}\otimes\vec{m}} = (\mathrm{id}_{\vec{l}} \otimes \beta^{\boldsymbol{\mathcal{C}}}_{\vec{k},\vec{m}}) \circ (\beta^{\boldsymbol{\mathcal{C}}}_{\vec{k},\vec{l}} \otimes \mathrm{id}_{\vec{m}}) \quad , \quad \beta^{\boldsymbol{\mathcal{C}}}_{\vec{k}\otimes\vec{l},\vec{m}} = (\beta^{\boldsymbol{\mathcal{C}}}_{\vec{k},\vec{m}} \otimes \mathrm{id}_{\vec{l}}) \circ (\mathrm{id}_{\vec{k}} \otimes \beta^{\boldsymbol{\mathcal{C}}}_{\vec{l},\vec{m}}). \tag{14}$$

Proposition 2.22. The pair $(\infty\text{-}\mathbf{Web}_{\mathrm{gr}}, \beta^{\bullet}_{\cdot,\cdot})$ is a braided monoidal category.

Proof. Since $\infty\text{-}\mathbf{Web}_{\mathrm{gr}}$ is a monoidal category and $\beta^{\bullet}_{\vec{k},\vec{l}}$ are isomorphism that clearly satisfy (14), we only need to prove that they are natural isomorphism. That is, we need to show that, for each web $u \in \mathrm{Hom}_{\infty\text{-}\mathbf{Web}_{\mathrm{gr}}}(\vec{k}, \vec{l})$ and each other object $\vec{m} = (m_1, \dots, m_c)$ of $\infty\text{-}\mathbf{Web}_{\mathrm{gr}}$, we have (we again use blue as a generic color)


m_1 ⋯ m_c l_1 ⋯ l_b
u
$\mathrm{id}_{\vec{m}}$
k_1 ⋯ k_a m_1 ⋯ m_c
=
m_1 ⋯ m_c l_1 ⋯ l_b
$\mathrm{id}_{\vec{m}}$
u
k_1 ⋯ k_a m_1 ⋯ m_c

The equality follows from Lemma 2.19. This proves the statement. □

The braiding $\beta^{\bullet}_{\cdot,\cdot}$ descends to the subquotients $N\text{-}\mathbf{Web}_{\mathrm{gr}}$, $N\text{-}\mathbf{Web}_{\mathrm{g}}$ and $N\text{-}\mathbf{Web}_{\mathrm{r}}$ and we denote all induced braidings also by $\beta^{\bullet}_{\cdot,\cdot}$. They are all given by the formulas in Definition 2.17, but some diagrams might be zero due to (11).

Corollary 2.23. $(N\text{-}\mathbf{Web}_{\mathrm{gr}}, \beta^{\bullet}_{\cdot,\cdot})$, $(N\text{-}\mathbf{Web}_{\mathrm{g}}, \beta^{\bullet}_{\cdot,\cdot})$ and $(N\text{-}\mathbf{Web}_{\mathrm{r}}, \beta^{\bullet}_{\cdot,\cdot})$, with $\beta^{\bullet}_{\cdot,\cdot}$ induced from $(\infty\text{-}\mathbf{Web}_{\mathrm{gr}}, \beta^{\bullet}_{\cdot,\cdot})$, are braided monoidal categories. □

Note that $N\text{-}\mathbf{Web}_{\mathrm{CKM}}$ is also a braided monoidal category, see [4, Corollary 6.2.3]. We rescale their braiding by multiplying it with $q^{\frac{kl}{N}}$ and we denote the resulting braided monoidal category by $(N\text{-}\mathbf{Web}_{\mathrm{CKM}}, \beta^{\bullet}_{\cdot,\cdot})$. The following corollary is immediate from Corollary 2.16.

Corollary 2.24. The functor $\iota_1^{\infty} \colon (N\text{-}\mathbf{Web}_{\mathrm{CKM}}, \beta^{\bullet}_{\cdot,\cdot}) \to (N\text{-}\mathbf{Web}_{\mathrm{gr}}, \beta^{\bullet}_{\cdot,\cdot})$ is an inclusion of a full, braided monoidal subcategory. □

2.5. A collection of diagrammatic idempotents.

Recall that the *Iwahori-Hecke algebra* $H_K(q)$ is the q-deformation of the symmetric group algebra $\mathbb{C}[S_K]$ in K letters. It is generated by $\{H_i \mid s_i \in S_K\}$ for all transpositions $s_i = (i, i+1) \in S_K$ subject to the relations

$$H_i^2 = (q - q^{-1})H_i + 1, \quad \text{for } i = 1, \dots, K-1,$$
$$H_iH_j = H_jH_i, \quad \text{for } |i-j| > 1 \quad , \quad H_iH_jH_i = H_jH_iH_j, \quad \text{for } |i-j| = 1.$$

There is a representation $p_K \colon \mathbb{C}_q(B_K) \to H_K(q)$ of the group algebra $\mathbb{C}_q(B_K)$ of the braid group B_K in K strands given by sending the braid group generators b_i (between the strands i and $i+1$) to H_i. Thinking of the generators H_i of $H_K(q)$ as crossings also makes sense from the perspective of the webs, as the next lemma shows.

Lemma 2.25. Given $K \in \mathbb{Z}_{\geq 0}$, then

$$\Phi_{q\mathrm{SW}}^{\infty}\colon H_K(q) \xrightarrow{\cong} \mathrm{End}_{\infty\text{-}\mathbf{Web}_{\mathrm{gr}}}(\vec{K}), \quad H_i \mapsto$$



$$, \quad \text{as } \mathbb{C}_q\text{-algebras.}$$

In order to prove Lemma 2.25, which will be used in Section 4, we need Theorem 3.20.

Proof. A direct computation shows that $\Phi_{q\mathrm{SW}}$ is a well-defined $\mathbb{C}_q$-algebra homomorphism. In fact, the composition $\Gamma \circ \Phi_{q\mathrm{SW}}^{\infty}$ is the isomorphism induced by quantum Schur-Weyl duality. To see this, let $V = (\mathbb{C}_q^N)^{\otimes K}$ and recall that quantum Schur-Weyl duality states that

$$\Phi_{q\mathrm{SW}}^{N}\colon H_K(q) \twoheadrightarrow \mathrm{End}_{\mathbf{U}_q(\mathfrak{gl}_N)}(V) \quad \text{and} \quad \Phi_{q\mathrm{SW}}^{N}\colon H_K(q) \xrightarrow{\cong} \mathrm{End}_{\mathbf{U}_q(\mathfrak{gl}_N)}(V), \text{ if } N \geq K. \tag{15}$$

Here $\Phi_{q\mathrm{SW}}^{N}$ is the $\mathbb{C}_q$-algebra homomorphism induced by the action of $H_K(q)$ on the K-fold tensor product V. By Theorem 3.20, we will get an isomorphism $H_K(q) \cong \mathrm{End}_{N\text{-}\mathbf{Web}_{\mathrm{gr}}}(\vec{K})$, if $N \geq K$. By using Proposition 2.15, there is a basis of $\mathrm{End}_{N\text{-}\mathbf{Web}_{\mathrm{gr}}}(\vec{K})$ for $N \geq K$ given by webs with only black edges or green edges with labels at most K. Since K is fixed, a direct comparison shows that $\Phi_{q\mathrm{SW}}^{\infty}$ has to be an isomorphism as well. □

Let $K \in \mathbb{Z}_{\geq 0}$ and let $\Lambda^+(K)$ denote the set of all *Young diagrams with K nodes*, e.g.

$\lambda = (4,3,1,1) \in \Lambda^+(9) \leftrightsquigarrow \lambda =$



$, \ \lambda^T = (4,2,2,1) \in \Lambda^+(9) \leftrightsquigarrow \lambda^{\mathrm{T}} =$



,

where we use the English notation for our Young diagrams. Here we have also displayed the *transpose* Young diagram λ^{T} of λ.

The following definition is motivated from [11] and [1].

Definition 2.26. (**Gyoja-Aiston idempotents**) Given $\lambda \in \Lambda^+(K)$, we associate to it a primitive *idempotent* $e_q(\lambda) \in \mathrm{End}_{\infty\text{-}\mathbf{Web}_{\mathrm{gr}}}(\vec{K})$. First we define two idempotents as tensor products of green or red clasps:

$$e_{\mathrm{col}}(\lambda) = \mathcal{CL}^{g}_{\mathrm{col}_1} \otimes \cdots \otimes \mathcal{CL}^{g}_{\mathrm{col}_c} \quad , \quad e_{\mathrm{row}}(\lambda) = \mathcal{CL}^{r}_{\mathrm{row}_1} \otimes \cdots \otimes \mathcal{CL}^{r}_{\mathrm{row}_r},$$

where c, r are the number of columns and rows of λ respectively, and col_i and row_i denote the number of boxes in the i-th column and row.

Denote by $T_\lambda^{\rightarrow}$ and by $T_\lambda^{\downarrow}$ the two tableaux of shape λ obtained by filling the numbers $1, \dots, K$ into Young diagram λ in order: $\rightarrow$ means rows before columns and $\downarrow$ means columns before rows (both from left to right). Pick any shortest permutation $w(\lambda) \in S_K$ permuting $T_\lambda^{\rightarrow}$ to $T_\lambda^{\downarrow}$. Then we define the *quasi-idempotent associated to* λ via

$$\tilde{e}_q(\lambda) = e_{\mathrm{col}}(\lambda) \circ \beta^{\bullet}_{w(\lambda)} \circ e_{\mathrm{row}}(\lambda) \circ \beta^{\bullet}_{w(\lambda)^{-1}}.$$

By [1, Theorem 4.7] (and the fact that their definition agrees with ours by Lemma 2.25 and Remark 2.21), there exists a non-zero scalar $a(\lambda) \in \mathbb{C}_q$ such that $\tilde{e}_q(\lambda)^2 = a(\lambda)\tilde{e}_q(\lambda)$. Thus, we define the *idempotent associated to* λ to be $e_q(\lambda) = \frac{1}{a(\lambda)}\tilde{e}_q(\lambda)$. These are primitive and orthogonal by [11, Theorem 4.5] and [1, Theorem 4.7]. ◇

Example 2.27. If $K = 2$, then there are two primitive idempotents, namely



Note that $a(\lambda) = 1$ for all one column or only one row Young diagrams λ. ■

Lemma 2.28. Exchanging green and red swaps $e_q(\lambda)$ to $\beta^{\bullet}_{w(\lambda)} e_q(\lambda^{\mathrm{T}}) \beta^{\bullet}_{w(\lambda)^{-1}}$.

Proof. An elementary computation, which uses that $e_{\mathrm{col}}(\lambda)$, $\beta^{\bullet}_{w(\lambda)} \circ e_{\mathrm{row}}(\lambda) \circ \beta^{\bullet}_{w(\lambda)^{-1}}$ and $e_q(\lambda)$ are idempotents and that $e_q(\lambda)$ is primitive, shows:

$$\tilde{e}_q(\lambda) = e_{\mathrm{col}}(\lambda) \circ \beta^{\bullet}_{w(\lambda)} \circ e_{\mathrm{row}}(\lambda) \circ \beta^{\bullet}_{w(\lambda)^{-1}} \circ e_{\mathrm{col}}(\lambda),$$
$$\beta^{\bullet}_{w(\lambda)} \tilde{e}_q(\lambda^{\mathrm{T}}) \beta^{\bullet}_{w(\lambda)^{-1}} = e_{\mathrm{row}}(\lambda^{\mathrm{T}}) \circ \beta^{\bullet}_{w(\lambda^{\mathrm{T}})^{-1}} \circ e_{\mathrm{col}}(\lambda^{\mathrm{T}}) \circ \beta^{\bullet}_{w(\lambda^{\mathrm{T}})} \circ e_{\mathrm{row}}(\lambda^{\mathrm{T}}).$$

Observe that $w(\lambda^{\mathrm{T}}) = w(\lambda)^{-1}$ and that $e_{\mathrm{col}}(\lambda)$ and $e_{\mathrm{row}}(\lambda)$ differ from $e_{\mathrm{row}}(\lambda^{\mathrm{T}})$ and $e_{\mathrm{col}}(\lambda^{\mathrm{T}})$ respectively only in exchanging the colors green and red. This proves the statement of the lemma for the quasi-idempotents. Applying the green-red symmetry to both sides of the equation $\tilde{e}_q(\lambda)^2 = a(\lambda)\tilde{e}_q(\lambda)$ shows that $a(\lambda) = a(\lambda^{\mathrm{T}})$ and the lemma follows. ☐

Example 2.29. For $\lambda = (3, 1) \in \Lambda^+(4)$, we have

$$\lambda = \begin{array}{|c|c|c|}\hline & & \\ \hline \\ \cline{1-1}\end{array} \quad , \quad T^{\rightarrow}_{\lambda} = \begin{array}{|c|c|c|}\hline 1 & 2 & 3 \\ \hline 4 \\ \cline{1-1}\end{array} \quad , \quad T^{\downarrow}_{\lambda} = \begin{array}{|c|c|c|}\hline 1 & 3 & 4 \\ \hline 2 \\ \cline{1-1}\end{array}.$$

Thus, $w = (243) \in S_4$ permutes $T^{\rightarrow}_{\lambda}$ to $T^{\downarrow}_{\lambda}$. Then


$\tilde{e}_q(\lambda) =$
1 1 1 1
$\mathcal{CL}^g_2$
$\beta^{\bullet}_{w(\lambda)}$
$\mathcal{CL}^r_3$
$\beta^{\bullet}_{w(\lambda)^{-1}}$
1 1 1 1
green↔red
1 1 1 1
$\mathcal{CL}^r_2$
$\beta^{\bullet}_{w(\lambda)}$
$\mathcal{CL}^g_3$
$\beta^{\bullet}_{w(\lambda)^{-1}}$
1 1 1 1
$\equiv$
1 1 1 1
$\mathcal{CL}^g_3$
$\beta^{\bullet}_{w(\lambda)^{-1}}$
$\mathcal{CL}^r_2$
$\beta^{\bullet}_{w(\lambda)}$
1 1 1 1
$= \tilde{e}_q(\lambda^{\mathrm{T}})$.

Moreover, the scaling factor in this case is $a(\lambda) = \frac{[4]}{[2][3]} = a(\lambda^{\mathrm{T}})$. ■

Remark 2.30. For $N \geq K$, the $H_K(q)$-module $(\mathbb{C}^N_q)^{\otimes K}$ decomposes into $\bigoplus_{\lambda \in \Lambda^+(K)} (S^{\lambda})^{\oplus m_{\lambda}}$ where the S^{λ} are the irreducible *Specht modules* for $H_K(q)$ and m_{λ} are their multiplicities. The primitive idempotents $e_q(\lambda)$ from Definition 2.26 are quantizations of Young symmetrizers that project onto S^{λ}. Note that a braid-conjugate of $e_q(\lambda)$, e.g. as in Lemma 2.28, might project onto a different copy of S^{λ} in the above decomposition.

3. PROOFS OF THE DIAGRAMMATIC PRESENTATIONS

This section contains the proof of our main theorems.

3.1. **Super q-Howe duality.**

Let $m,n\in\mathbb{Z}_{\geq 0}$. We start by recalling the *quantum general linear superalgebra* $\mathbf{U}_q(\mathfrak{gl}_{m|n})$ and introduce its *idempotented form* $\dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n})$. We follow the conventions used in [31], but adapt Zhang's notation to be closer to the one from [4].

To this end, recall that the $\mathfrak{gl}_{m|n}$-weight lattice is isomorphic to $\mathbb{Z}^{m+n}$ and we denote the $\mathfrak{gl}_{m|n}$-weights usually by vectors $\vec{k}=(k_1,\dots,k_m,k_{m+1},\dots,k_{m+n})$. For $\mathbb{I}=\mathbb{I}_0\cup\mathbb{I}_1$ define

$$|i|=\begin{cases}0, & \text{if } i\in\mathbb{I}_0=\{1,\dots,m\},\\ 1, & \text{if } i\in\mathbb{I}_1=\{m+1,\dots,m+n\}.\end{cases}$$

The notation $|\cdot|$ means the *super* degree (which is a $\mathbb{Z}/2$-degree). We use a similar notation for all $\mathbb{Z}/2$-graded spaces where we, by convention, always consider degrees modulo 2 in the following. Moreover, let $\epsilon_i=(0,\dots,0,1,0,\dots,0)\in\mathbb{Z}^{m+n}$, with 1 being in the i-th coordinate, and denote by $\alpha_i=\epsilon_i-\epsilon_{i+1}=(0,\dots,1,-1,\dots,0)\in\mathbb{Z}^{m+n}$ for $i\in\mathbb{I}-\{m+n\}$ the i-th simple root. Recall that the *super* Euclidean inner product on $\mathbb{Z}^{m+n}$ is given by $(\epsilon_i,\epsilon_j)_{\mathrm{su}}=(-1)^{|i|}\delta_{i,j}$.

Definition 3.1. Let $m,n\in\mathbb{Z}_{\geq 0}$. The *quantum general linear superalgebra* $\mathbf{U}_q(\mathfrak{gl}_{m|n})$ is the associative, $\mathbb{Z}/2$-graded, unital $\mathbb{C}_q$-algebra generated by $L_i^{\pm 1}$, for $i\in\mathbb{I}$, and F_i,E_i, for $i\in\mathbb{I}-\{m+n\}$, subject to the *non-super* relations

$$L_iL_j=L_jL_i,\quad L_iL_i^{-1}=L_i^{-1}L_i=1,\quad L_iF_j=q^{-(\epsilon_i,\alpha_j)_{\mathrm{su}}}F_jL_i,\quad L_iE_j=q^{(\epsilon_i,\alpha_j)_{\mathrm{su}}}E_jL_i,$$

$$E_iF_j-F_jE_i=\delta_{i,j}\frac{L_iL_{i+1}^{-1}-L_i^{-1}L_{i+1}}{q-q^{-1}},\quad i\neq m,$$

$$[2]F_iF_jF_i=F_i^2F_j+F_jF_i^2,\quad \text{if } |i-j|=1,\ i\neq m,\qquad F_iF_j-F_jF_i=0,\quad \text{if } |i-j|>1,$$

$$[2]E_iE_jE_i=E_i^2E_j+E_jE_i^2,\quad \text{if } |i-j|=1,\ i\neq m,\qquad E_iE_j-E_jE_i=0,\quad \text{if } |i-j|>1,$$

(for suitable $i,j\in\mathbb{I}$) and the *super relations*

$$F_m^2=0=E_m^2,\quad E_mF_m+F_mE_m=\frac{L_mL_{m+1}^{-1}-L_m^{-1}L_{m+1}}{q-q^{-1}},$$

$$\begin{aligned}[2]F_mF_{m+1}F_{m-1}F_m ={}& F_mF_{m+1}F_mF_{m-1}+F_{m-1}F_mF_{m+1}F_m\\ &+F_{m+1}F_mF_{m-1}F_m+F_mF_{m-1}F_mF_{m+1},\\ [2]E_mE_{m+1}E_{m-1}E_m ={}& E_mE_{m+1}E_mE_{m-1}+E_{m-1}E_mE_{m+1}E_m\\ &+E_{m+1}E_mE_{m-1}E_m+E_mE_{m-1}E_mE_{m+1}.\end{aligned}$$

Moreover, $|L_i|=0$ for $i\in\mathbb{I}$, $|F_i|=|E_i|=0$ for $i\in\mathbb{I}-\{m\}$ and $|F_m|=|E_m|=1$. ◇

We recover $\mathbf{U}_q(\mathfrak{gl}_N)$ by setting $m=N$ and $n=0$. We denote $\mathbb{I}_N=\{1,\dots,N\}$ in the following to distinguish it from $\mathbb{I}$ as above. Note that $\mathbf{U}_q(\mathfrak{gl}_N)$ is concentrated in degree 0.

The algebra $\mathbf{U}_q(\mathfrak{gl}_{m|n})$ is a $\mathbb{Z}/2$-graded Hopf algebra with coproduct Δ, antipode S and the counit ε given by

$$\Delta(F_i)=F_i\otimes 1+L_i^{-1}L_{i+1}\otimes F_i,\quad \Delta(E_i)=E_i\otimes L_iL_{i+1}^{-1}+1\otimes E_i,\quad \Delta(L_i)=L_i\otimes L_i,$$

$$S(F_i)=-L_iL_{i+1}^{-1}F_i,\ S(E_i)=-E_iL_i^{-1}L_{i+1},\ S(L_i)=L_i^{-1},\quad \varepsilon(F_i)=\varepsilon(E_i)=0,\ \varepsilon(L_i)=1.$$

In the spirit of Beilinson, Lusztig and MacPherson [2], we now adjoin for all $\vec{k} \in \mathbb{Z}^{m+n}$ idempotents $1_{\vec{k}}$ of super degree $|1_{\vec{k}}| = 0$ to $\mathbf{U}_q(\mathfrak{gl}_{m|n})$. Denote by I the ideal generated by the relations

$$\begin{aligned} 1_{\vec{k}}1_{\vec{l}} = \delta_{\vec{k},\vec{l}}1_{\vec{k}}, &\qquad L_i 1_{\vec{k}} = q^{k_i(\epsilon_i,\epsilon_i)_{\mathrm{su}}}1_{\vec{k}}, \\ 1_{\vec{k}-\alpha_i}F_i1_{\vec{k}} = F_i1_{\vec{k}} = 1_{\vec{k}-\alpha_i}F_i, &\qquad 1_{\vec{k}+\alpha_i}E_i1_{\vec{k}} = E_i1_{\vec{k}} = 1_{\vec{k}+\alpha_i}E_i. \end{aligned}$$

Definition 3.2. Define by

$$\dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n}) = \bigoplus_{\vec{k},\vec{l}\in\mathbb{Z}^{m+n}} 1_{\vec{l}}\mathbf{U}_q(\mathfrak{gl}_{m|n})1_{\vec{k}} \mod I$$

the *idempotented* quantum general linear superalgebra. ◇

Remark 3.3. One can view $\dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n})$ as generated by the *divided powers*

$$F_i^{(j)} = \frac{F_i^j}{[j]!} \quad \text{and} \quad E_i^{(j)} = \frac{E_i^j}{[j]!}, \quad \text{for } i \in \mathbb{I}{-}\{m+n\}.$$

This allows the definition of an integral version of $\dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n})$. For simplicity, we work over $\mathbb{C}_q$ in this paper and we do not consider the integral version.

The relations in $\dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n})$ are obtained from the relations of $\mathbf{U}_q(\mathfrak{gl}_{m|n})$. For convenience we list the new versions of the *super* relations:

$$\begin{aligned} F_m^2 1_{\vec{k}} = 0 = E_m^2 1_{\vec{k}}, &\qquad E_mF_m1_{\vec{k}} + F_mE_m1_{\vec{k}} = [k_m + k_{m+1}]1_{\vec{k}}, \\ [2]F_mF_{m+1}F_{m-1}F_m1_{\vec{k}} =& F_mF_{m+1}F_mF_{m-1}1_{\vec{k}} + F_{m-1}F_mF_{m+1}F_m1_{\vec{k}} \\ &+ F_{m+1}F_mF_{m-1}F_m1_{\vec{k}} + F_mF_{m-1}F_mF_{m+1}1_{\vec{k}}, \end{aligned} \tag{16}$$

the second of which we call the *super commutation relation* (the third type of relation holds for E's as well).

It is convenient for us hereinafter to view $\dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n})$ as a category whose objects are the $\mathfrak{gl}_{m|n}$-weights $\vec{k} \in \mathbb{Z}^{m+n}$ and $\mathrm{Hom}_{\dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n})}(\vec{k},\vec{l}) = 1_{\vec{l}}\dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n})1_{\vec{k}}$.

Recall that the vector representation $\mathbb{C}_q^{m|n}$ of $\mathbf{U}_q(\mathfrak{gl}_{m|n})$ has a basis given by $\{x_i \mid i \in \mathbb{I}\}$ with super degrees $|x_i| = |i|$ for $i \in \mathbb{I}$, where the $\mathbf{U}_q(\mathfrak{gl}_{m|n})$-action is defined via

$$F_i(x_j) = \begin{cases} x_{j+1}, & \text{if } i = j, \\ 0, & \text{else,} \end{cases} \quad E_i(x_j) = \begin{cases} x_{j-1}, & \text{if } i = j-1, \\ 0, & \text{else,} \end{cases} \quad L_i(x_j) = q^{(\epsilon_i,\epsilon_j)_{\mathrm{su}}}x_j.$$

We need to consider the *quantum exterior superalgebra* $\bigwedge_q^{\bullet}(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^N)$. Recall that a vector space $V = V_0 \oplus V_1$ with a $\mathbb{Z}/2$-grading is called a *super* vector space. Here V_0 and V_1 are its degree 0 and 1 parts. These graded parts of $\mathbb{C}_q^{m|n}$ have bases given by $\{x_i \mid i \in \mathbb{I}_0\}$ and $\{x_i \mid i \in \mathbb{I}_1\}$ respectively. In contrast, $\mathbb{C}_q^N = (\mathbb{C}_q^N)_0$ is concentrated in degree zero and we denote its basis by $\{y_j \mid j \in \mathbb{I}_N\}$. Additionally, the tensor product $V \otimes W$ of two super vector spaces V and W is a super vector space with $v \otimes w$ of degree $|v| + |w|$ for two homogeneous elements v, w. Specifically, $\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^N$ is a super vector space with $(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^N)_0$ spanned by $\{z_{ij} = x_i \otimes y_j \mid i \in \mathbb{I}_0,\ j \in \mathbb{I}_N\}$ and $(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^N)_1$ spanned by $\{z_{ij} = x_i \otimes y_j \mid i \in \mathbb{I}_1,\ j \in \mathbb{I}_N\}$.

Here $|z_{ij}| = |i|$. Note that $(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^N)^{\otimes K}$ is a $\mathbb{Z}/2$-graded $\mathbf{U}_q(\mathfrak{gl}_{m|n}) \otimes \mathbf{U}_q(\mathfrak{gl}_N)$-module for all $K \in \mathbb{Z}_{\geq 0}$ by using the Hopf algebras structures of $\mathbf{U}_q(\mathfrak{gl}_{m|n})$ and $\mathbf{U}_q(\mathfrak{gl}_N)$.

We denote by $\mathrm{Sym}_q^2(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^N)$ the *second symmetric super power* as in [21, (4.1)]. Armed with this notation, we define the *quantum exterior superalgebra*

$$\textstyle\bigwedge_q^\bullet(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^N) = T(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^N)/\mathrm{Sym}_q^2(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^N),$$

where $T(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^N) = \bigoplus_{K \in \mathbb{Z}_{\geq 0}} (\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^N)^{\otimes K}$ denotes the *super tensor algebra* of $\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^N$. This is a $\mathbf{U}_q(\mathfrak{gl}_{m|n}) \otimes \mathbf{U}_q(\mathfrak{gl}_N)$-module and decompose as

$$\textstyle\bigwedge_q^\bullet(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^N) \cong \displaystyle\bigoplus_{K \in \mathbb{Z}_{\geq 0}} \textstyle\bigwedge_q^K(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^N).$$

The space $\bigwedge_q^K(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^N)$ is called the *degree K part* of $\bigwedge_q^\bullet(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^N)$.

Remark 3.4. We recover the degree K part of the quantum exterior algebra $\bigwedge_q^K(\mathbb{C}_q^m \otimes \mathbb{C}_q^N)$ by setting $n = 0$ and, by [25, Remark 2.1], the degree K part of the quantum symmetric algebra $\mathrm{Sym}_q^K(\mathbb{C}_q^n \otimes \mathbb{C}_q^N)$ by setting $m = 0$. These algebras were originally defined in [3, Definition 2.7] and used in [4, Section 4.2] and in [23, Section 2.1] respectively to study skew and symmetric q-Howe duality.

Example 3.5. $\bigwedge_q^K(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^N)$ has a basis given by

$$\left\{ z_{i_1 j_1} \otimes \cdots \otimes z_{i_K j_K} \middle| \begin{array}{l} 1 \leq i_1 \leq \cdots \leq i_K \leq m+n, \\ 1 \leq j_1 \leq \cdots \leq j_K \leq N, \end{array} \text{ and } \begin{array}{l} i_k = i_{k+1}, \\ j_k = j_{k+1}, \end{array} \Rightarrow |i_k| = 1 \right\}. \tag{17}$$

This can be deduced from [21, Lemma 4.1]. In particular, by setting $m = 1$ and $n = 0$, we obtain a basis for $\bigwedge_q^K \mathbb{C}_q^N$ of the form

$$\{ y_{i_1} \otimes \cdots \otimes y_{i_K} \mid 1 \leq y_1 < \cdots < y_K \leq N \}, \tag{18}$$

while setting $m = 0$ and $n = 1$ gives a basis for $\mathrm{Sym}_q^K \mathbb{C}_q^N$ of the form

$$\{ y_{i_1} \otimes \cdots \otimes y_{i_K} \mid 1 \leq y_1 \leq \cdots \leq y_K \leq N \}. \tag{19}$$

These are precisely the usual (non-super) bases, see for example [3, Section 2.4]. ■

We call a $\mathfrak{gl}_{m|n}$-weight $\lambda = (\lambda_1, \ldots, \lambda_{m+n}) \in \mathbb{Z}^{m+n}$ a *dominant integral* $\mathfrak{gl}_{m|n}$-weight if it is dominant integral as a $\mathfrak{gl}_m \oplus \mathfrak{gl}_n$-weight. We need these only in the case where they can be seen as $(m|n)$-hook Young diagrams that we, abusing notation, also denote by λ (here we use the conventions for these $(m|n)$-hook Young diagrams as in [5, Theorem 3.3]). Moreover, we call a dominant integral $\mathfrak{gl}_{m|n}$-weight λ an *$(m|n, N)$-supported $\mathfrak{gl}_{m|n}$-weight* if it corresponds to an $(m|n)$-hook Young diagram with at most N columns. For each such λ there exists an irreducible $\mathbf{U}_q(\mathfrak{gl}_{m|n})$-module $L_{m|n}(\lambda)$ respectively $\mathbf{U}_q(\mathfrak{gl}_N)$-module $L_N(\lambda^{\mathrm{T}})$, see for example [16, Section 2.5].

Theorem 3.6. (**Super q-Howe duality**) We have the following.

(a) Let $K \in \mathbb{Z}_{\geq 0}$. The actions of $\mathbf{U}_q(\mathfrak{gl}_{m|n})$ and $\mathbf{U}_q(\mathfrak{gl}_N)$ on $\bigwedge_q^K(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^N)$ commute and generate each others commutant.

(b) There exists an isomorphism

$$\textstyle\bigwedge_q^\bullet(\mathbb{C}_q^{m|n}\otimes\mathbb{C}_q^N)\cong(\bigwedge_q^\bullet\mathbb{C}_q^N)^{\otimes m}\otimes(\mathrm{Sym}_q^\bullet\mathbb{C}_q^N)^{\otimes n}$$

of $\mathbf{U}_q(\mathfrak{gl}_N)$-modules under which the $\vec{k}$-weight space of $\bigwedge_q^\bullet(\mathbb{C}_q^{m|n}\otimes\mathbb{C}_q^N)$ (considered as a $\mathbf{U}_q(\mathfrak{gl}_{m|n})$-module) is identified with

$$\textstyle\bigwedge_q^{\vec{k}_0}\mathbb{C}_q^N\otimes\mathrm{Sym}_q^{\vec{k}_1}\mathbb{C}_q^N=\bigwedge_q^{k_1}\mathbb{C}_q^N\otimes\cdots\otimes\bigwedge_q^{k_m}\mathbb{C}_q^N\otimes\mathrm{Sym}_q^{k_{m+1}}\mathbb{C}_q^N\otimes\cdots\otimes\mathrm{Sym}_q^{k_{m+n}}\mathbb{C}_q^N. \tag{20}$$

Here $\vec{k}=(k_1,\dots,k_{m+n})$, $\vec{k}_0=(k_1,\dots,k_m)$ and $\vec{k}_1=(k_{m+1},\dots,k_{m+n})$.

(c) As $\mathbf{U}_q(\mathfrak{gl}_{m|n})\otimes\mathbf{U}_q(\mathfrak{gl}_N)$-modules, we have a decomposition of the form

$$\textstyle\bigwedge_q^K(\mathbb{C}_q^{m|n}\otimes\mathbb{C}_q^N)\cong\bigoplus_\lambda L_{m|n}(\lambda)\otimes L_N(\lambda^{\mathrm{T}}),$$

where we sum over all $(m|n,N)$-supported $\mathfrak{gl}_{m|n}$-weights λ whose entries sum up to K. This induces a decomposition

$$\textstyle\bigwedge_q^\bullet(\mathbb{C}_q^{m|n}\otimes\mathbb{C}_q^n)\cong\bigoplus_\lambda L_{m|n}(\lambda)\otimes L_N(\lambda^{\mathrm{T}}),$$

where we sum over all $(m|n,N)$-supported $\mathfrak{gl}_{m|n}$-weights λ.

Remark 3.7. Symmetric and skew Howe duality for the pair $(\mathrm{GL}_m,\mathrm{GL}_N)$ is originally due to Howe, see [12, Section 2 and Section 4]. Note that the non-quantum version of Theorem 3.6 can be found for example in [5, Theorem 3.3] or [25, Proposition 2.2]. Moreover, the "dual" of Theorem 3.6, given by considering $\mathbf{U}_q(\mathfrak{gl}_N)$ as the Howe dual group instead of $\mathbf{U}_q(\mathfrak{gl}_{m|n})$, can be found in [21, Proposition 4.3].

Proof. (a) and (c) are proven in [28, Theorem 2.2] or in [21, Theorem 4.2] and only (b) remains to be verified. For this purpose, we use the bases from (17), (18) and (19) to define:

$$\textstyle T_i^e\colon\bigwedge_q^k(\mathbb{C}_q^N)\to\bigwedge_q^k(\mathbb{C}_q^{m|n}\otimes\mathbb{C}_q^N),\quad y_{j_1}\otimes\cdots\otimes y_{j_k}\mapsto z_{ij_1}\otimes\cdots\otimes z_{ij_k},\quad i\in\mathbb{I}_0,$$

$$\textstyle T_i^s\colon\mathrm{Sym}_q^k(\mathbb{C}_q^n)\to\bigwedge_q^k(\mathbb{C}_q^{m|n}\otimes\mathbb{C}_q^N),\quad y_{j_1}\otimes\cdots\otimes y_{j_k}\mapsto z_{ij_1}\otimes\cdots\otimes z_{ij_k},\quad i\in\mathbb{I}_1.$$

That these maps are well-defined $\mathbf{U}_q(\mathfrak{gl}_N)$-intertwiners follows from the explicit description in Example 3.5. Injectivity was shown in [4, Theorem 4.2.2] for the first and in [23, Theorem 2.6] for the second map. Thus, for $\vec{k}\in\mathbb{Z}^{m+n}$ with $k_1+\cdots+k_{m+n}=K$, we see that

$$\textstyle T\colon\bigoplus_{\vec{k}\in\mathbb{Z}_{\geq 0}^{m+n}}\bigwedge_q^{\vec{k}_0}\mathbb{C}_q^N\otimes\mathrm{Sym}_q^{\vec{k}_1}\mathbb{C}_q^N\to\bigwedge_q^K(\mathbb{C}_q^{m|n}\otimes\mathbb{C}_q^N)$$

given by $T(v_1\otimes\cdots\otimes v_{m+n})=T_1^e(v_1)\otimes\cdots\otimes T_m^e(v_m)\otimes T_{m+1}^s(v_{m+1})\otimes\cdots\otimes T_{m+n}^s(v_{m+n})$, is an $\mathbf{U}_q(\mathfrak{gl}_N)$-module isomorphism by comparing the sizes of the bases from Example 3.5. This clearly induces the isomorphism of $\mathbf{U}_q(\mathfrak{gl}_N)$-modules we are looking for.

It remains to verify the $\mathbf{U}_q(\mathfrak{gl}_{m|n})$-weight space decomposition from (20). To this end, we only have to see that the action on $\bigwedge_q^{\vec{k}_0}\mathbb{C}_q^N\otimes\mathrm{Sym}_q^{\vec{k}_1}\mathbb{C}_q^N$ of the $L_{i'}$'s of $\mathbf{U}_q(\mathfrak{gl}_{m|n})$ under the inverse of T is just a multiplication with $q^{k_i(\epsilon_i,\epsilon_{i'})_{\mathrm{su}}}$. The action of $\mathbf{U}_q(\mathfrak{gl}_{m|n})$ is given by

$$L_{i'}(z_{ij_1}\otimes\cdots\otimes z_{ij_{m+n}})=L_{i'}(z_{ij_1})\otimes\cdots\otimes L_{i'}(z_{ij_{m+n}})=q^{k_i(\epsilon_i,\epsilon_{i'})_{\mathrm{su}}}z_{ij_1}\otimes\cdots\otimes z_{ij_{m+n}}.$$

Hence, the $\mathbf{U}_q(\mathfrak{gl}_{m|n})$-weight space decomposition follows. □

By Theorem 3.6 part (b): we get linear maps

$$f_{\vec{k}}^{\vec{l}}\colon 1_{\vec{l}}\dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n})1_{\vec{k}} \to \mathrm{Hom}_{\mathbf{U}_q(\mathfrak{gl}_N)}(\textstyle\bigwedge_q^{\vec{k}_0}\mathbb{C}_q^N \otimes \mathrm{Sym}_q^{\vec{k}_1}\mathbb{C}_q^N, \textstyle\bigwedge_q^{\vec{l}_0}\mathbb{C}_q^N \otimes \mathrm{Sym}_q^{\vec{l}_1}\mathbb{C}_q^N)$$

for any two $\vec{k},\vec{l} \in \mathbb{Z}_{\geq 0}^{m+n}$ such that $\sum_{i=0}^{m+n} k_i = \sum_{i=0}^{m+n} l_i$. Using part (a) of Theorem 3.6, we see that the homomorphisms $f_{\vec{k}}^{\vec{l}}$ are all surjective. Thus, we get the following.

Corollary 3.8. There exists a full functor $\Phi_{\mathrm{su}}^{m|n}\colon \dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n}) \to \mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}$, which we call the *super q-Howe functor*, given on objects and morphisms by

$$\vec{k} \overset{\Phi_{\mathrm{su}}^{m|n}}{\longmapsto} \textstyle\bigwedge_q^{\vec{k}_0}\mathbb{C}_q^N \otimes \mathrm{Sym}_q^{\vec{k}_1}\mathbb{C}_q^N \quad , \quad 1_{\vec{l}}x1_{\vec{k}} \overset{\Phi_{\mathrm{su}}^{m|n}}{\longmapsto} f_{\vec{k}}^{\vec{l}}(x).$$

Everything else is sent to zero. □

3.2. **The sorted equivalences.** In this subsection we construct a full and faithful functor

$$\Gamma_N^{\mathrm{sort}}\colon N\text{-}\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}} \to \mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}^{\mathrm{sort}},$$

where $N\text{-}\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}}$ is the sorted web category from Definition 2.5 and $\mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}^{\mathrm{sort}}$ denotes the full subcategory of $\mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}$ whose objects are sorted as in (20).

We essentially define Γ_N^{sort} such that there is a commuting diagram

$$\begin{tikzcd} \dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n}) \arrow[r, "\Phi_{\mathrm{su}}^{m|n}"] \arrow[dr, "\Upsilon_{\mathrm{su}}^{m|n}"'] & \mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}^{\mathrm{sort}} \\ & N\text{-}\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}} \arrow[u, "\Gamma_N^{\mathrm{sort}}"'] \end{tikzcd} \tag{21}$$

The functor $\Upsilon_{\mathrm{su}}^{m|n}$ is a *ladder functor*, whose definition is motivated by [4, Subsection 5.1].

Lemma 3.9. Let $m,n \in \mathbb{Z}_{\geq 0}$. There exists a functor $\Upsilon_{\mathrm{su}}^{m|n}\colon \dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n}) \to N\text{-}\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}}$ which sends a $\mathfrak{gl}_{m|n}$-weight $\vec{k} \in \mathbb{Z}_{\geq 0}^{m+n}$ to $((k_1)_g,\dots,(k_m)_g,(k_{m+1})_r,\dots,(k_{m+n})_r)$ in $N\text{-}\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}}$ and all other $\mathfrak{gl}_{m|n}$-weights to the zero object. On morphisms $\Upsilon_{\mathrm{su}}^{m|n}$ is given by

$F_i^{(j)}1_{\vec{k}} \mapsto$ 

, $F_i^{(j)}1_{\vec{k}} \mapsto$ 

for $i \in \mathbb{I}_0 - \{m\}$ or $i \in \mathbb{I}_1 - \{m+n\}$ respectively, and

$F_m 1_{\vec{k}} \mapsto$ 

Similarly, but with reversed horizontal orientations, for the generators $E_i^{(j)}1_{\vec{k}}$ and $E_m 1_{\vec{k}}$.

Proof. To show that $\Upsilon_{\mathrm{su}}^{m|n}$ is well-defined, it suffices to show that all relations in $\dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n})$ are satisfied in N-$\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}}$. For monochromatic relations we can copy [4, Proposition 5.2.1]. Lemma 2.10 shows that the super relations (16) hold in N-$\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}}$. $\square$

Definition 3.10. (**The diagrammatic presentation functor** Γ_N^{sort}) We define a functor $\Gamma_N^{\mathrm{sort}}\colon N\text{-}\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}} \to \mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}^{\mathrm{sort}}$ as follows.

- On objects: to each $\vec{k} = ((k_1)_g, \dots, (k_m)_g, (k_{m+1})_r, \dots, (k_{m+n})_r)$, we assign
$$\Gamma_N^{\mathrm{sort}}(\vec{k}) = {\textstyle\bigwedge}_q^{\vec{k}_0}\mathbb{C}_q^N \otimes \mathrm{Sym}_q^{\vec{k}_1}\mathbb{C}_q^N,$$
where $\vec{k}_0 = (k_1, \dots, k_m)$ and $\vec{k}_1 = (k_{m+1}, \dots, k_{m+n})$. Moreover, we send the empty tuple to the trivial $\mathbf{U}_q(\mathfrak{gl}_N)$-module $\mathbb{C}_q$ and the zero object to the $\mathbf{U}_q(\mathfrak{gl}_N)$-module 0.
- On morphisms: we use the functor $\Phi_{\mathrm{su}}^{m|n}$ from Corollary 3.8 to define Γ_N^{sort} on the generating trivalent vertices in N-$\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}}$ (here we assume that the diagrams are the identities outside of the illustrated part). For this, let $i \in \mathbb{I}$ and we use the notation $k = k_i, l = k_{i+1}$ and $(k,l) = (k_1, \dots, k_i = k, k_{i+1} = l, \dots, k_{m+n})$.

(22)

Γ_N^{sort} () $= \Phi_{\mathrm{su}}^{m|n}(E_i^{(l)} 1_{(k,l)})$, Γ_N^{sort} () $= \Phi_{\mathrm{su}}^{m|n}(F_i^{(l)} 1_{(k+l,0)})$

Γ_N^{sort} () $= \Phi_{\mathrm{su}}^{m|n}(F_i^{(k)} 1_{(k,l)})$, Γ_N^{sort} () $= \Phi_{\mathrm{su}}^{m|n}(E_i^{(k)} 1_{(0,k+l)})$.

Note that these definitions include the mixed case, where we either have $l = 1$ (and colored black) or $k = 1$ (and colored black) and we use the odd generators F_m and E_m. $\diamond$

Remark 3.11. There are certain choices for the images of monochromatic merges and splits, but these choices do not matter, see [23, Remark 2.18]. In contrast, there is no other choice for the mixed merges and splits. For example, take $l = 1$ in the top left in (22). The green edge labeled $k+1$ should represent ${\textstyle\bigwedge}_q^{k+1}\mathbb{C}_q^N$. Thus, we have to see the top boundary of the left-hand side as $1_{(k+1,0)}$ and not as $1_{(0,k+1)}$, which determines our choices. Similarly for the other mixed generators. For example, if $m = n = 1$, and $k = 1$ or $l = 1$, then

Γ_N^{sort} () $= \Phi_{\mathrm{su}}^{1|1}(E_1 1_{(1,1)}) \neq \Phi_{\mathrm{su}}^{1|1}(F_1 1_{(1,1)}) = \Gamma_N^{\mathrm{sort}}$ ().

Lemma 3.12. Γ_N^{sort} is a well-defined functor $\Gamma_N^{\mathrm{sort}}\colon N\text{-}\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}} \to \mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}^{\mathrm{sort}}$ making the diagram (21) commutative.

Proof. First we note that $\Gamma_N^{\mathrm{sort}} \circ \Upsilon_{\mathrm{su}}^{m|n} = \Phi_{\mathrm{su}}^{m|n}$ on generators $F_i^{(j)} 1_{\vec{k}}$ and $F_m 1_{\vec{k}}$ (and analogously for E's) with $i \in \mathbb{I} - \{m\}, j \in \mathbb{Z}_{\geq 0}$ and $\vec{k} \in \mathbb{Z}^{m+n}$. This follows from the definition of Γ_N^{sort} via $\Phi_{\mathrm{su}}^{m|n}$ and the observation that ladders can be written as compositions of merges and splits, see also [23, Lemma 2.20].

We need to check that the images of the relations from N-$\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}}$ under Γ_N^{sort} hold in $\mathfrak{gl}_N$-$\mathbf{Mod}_{\mathrm{es}}^{\mathrm{sort}}$. Corollary 3.8 guarantees that all relations in $\mathfrak{gl}_N$-$\mathbf{Mod}_{\mathrm{es}}^{\mathrm{sort}}$ are induced via $\Phi_{\mathrm{su}}^{m|n}$ from relations in $\dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n})$ and the fact that $\Phi_{\mathrm{su}}^{m|n}$ kills certain $\mathfrak{gl}_{m|n}$-weights. It remains to check that the relations in N-$\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}}$ are, likewise, induced via $\Upsilon_{\mathrm{su}}^{m|n}$ from relations in $\dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n})$. For the monochromatic and isotopy relations, this follows as in [23, Lemma 2.20].

The dumbbell relation (10) can be recovered from $\dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n})$ as follows. Without loss of generality we work with $m = n = 1$:

$[2]$ 

$= \Upsilon_{\mathrm{su}}^{1|1}([2]1_{(1,1)}) = \Upsilon_{\mathrm{su}}^{1|1}(FE1_{(1,1)} + EF1_{(1,1)}) =$ 

$+$ 

Relation (11) is a consequence of killing $\mathfrak{gl}_{m|n}$-weights $\vec{k} = (k_1, \dots, k_{m+n})$, one of whose first m entries is larger than N. ☐

Lemma 3.13. The functor $\iota_1^\infty \colon N\text{-}\mathbf{Web}_{\mathrm{CKM}} \to N\text{-}\mathbf{Web}_{\mathrm{gr}}$ is faithful.

Proof. By Lemma 3.12 and a comparison of definitions, we have a commuting diagram

$$\begin{tikzcd}
\mathfrak{gl}_N\text{-}\mathbf{Mod}_e \arrow[r, "\iota_{\mathrm{e}}^{\mathrm{es}}"] & \mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}^{\mathrm{sort}} \\
N\text{-}\mathbf{Web}_{\mathrm{CKM}} \arrow[u, "\Gamma_{\mathrm{CKM}}"] \arrow[r, "\iota_1^\infty"'] & N\text{-}\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}} \arrow[u, "\Gamma_N^{\mathrm{sort}}"']
\end{tikzcd},$$

where Γ_{CKM} is the functor considered in [4, Subsection 3.2] and $\iota_{\mathrm{e}}^{\mathrm{es}}$ is the evident embedding of a full subcategory. Γ_{CKM} is faithful by [4, Theorem 3.3.1] and thus, ι_1^∞ is faithful as well. ☐

Remark 3.14. Let $\mathbf{Mat}(N\text{-}\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}})$ be the *additive closure* of N-$\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}}$: objects are finite, formal direct sums of the objects of N-$\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}}$ and morphisms are matrices (whose entries are morphisms from N-$\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}}$). We can extend Γ_N^{sort} additively to a functor

$$\Gamma_N^{\mathrm{sort}} \colon \mathbf{Mat}(N\text{-}\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}}) \to \mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}^{\mathrm{sort}}.$$

Similarly for Γ_N later on.

Proposition 3.15. The functor $\Gamma_N^{\mathrm{sort}} \colon N\text{-}\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}} \to \mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}^{\mathrm{sort}}$ gives rise to an equivalence of categories $\Gamma_N^{\mathrm{sort}} \colon \mathbf{Mat}(N\text{-}\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}}) \to \mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}^{\mathrm{sort}}$.

Proof. Since, by Lemma 3.12 and Remark 3.14, $\Gamma_N^{\mathrm{sort}} \colon \mathbf{Mat}(N\text{-}\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}}) \to \mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}^{\mathrm{sort}}$ is well-defined, it remains to show that Γ_N^{sort} is essentially surjective, full and faithful.

Essentially surjective: this follows directly from the definitions of Γ_N^{sort}, N-$\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}}$, its additive closure $\mathbf{Mat}(N\text{-}\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}})$ and $\mathfrak{gl}_N$-$\mathbf{Mod}_{\mathrm{es}}^{\mathrm{sort}}$.

Full: it suffices to verify fullness for morphisms between objects of the form $\vec{k} \in X^{m+n}$, where $X^{m+n} = (X_b \cup X_g)^m \cup (X_b \cup X_r)^n$. That it holds is clear from diagram (21) since $\Phi_{\mathrm{su}}^{m|n}$ is full by Corollary 3.8.

Faithful: again it suffices to verify faithfulness for morphisms between objects of the form $\vec{k} \in X^{m+n}$. Given any web $u \in \mathrm{Hom}_{N\text{-}\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}}}(\vec{k}, \vec{l})$ for $\vec{k} \in X^{m+n}$ and $\vec{l} \in X^{m'+n'}$, we can compose u from the bottom and the top with merges and splits respectively to obtain



Recall that exploding edges is, by (8), a reversible operation. Hence, we have

$$\Gamma_N^{\mathrm{sort}}(u) = \Gamma_N^{\mathrm{sort}}(v) \quad \text{iff} \quad \Gamma_N^{\mathrm{sort}}(u') = \Gamma_N^{\mathrm{sort}}(v'),$$

which together with Corollary 2.16 reduces the verification of faithfulness to the case where all web edges are black or green. Such webs lie in $\iota_1^{\infty}(N\text{-}\mathbf{Web}_{\mathrm{CKM}})$ and faithfulness follows as in the proof of Lemma 3.13. □

3.3. Proofs of the equivalences.

Remark 3.16. Recall that the *universal R-matrix* for $\mathfrak{gl}_N$ gives a braiding on $\mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}$ as follows (see [27, Chapter XI, Sections 2 and 7] and the references therein). For any pair of $\mathbf{U}_q(\mathfrak{gl}_N)$-modules V, W in $\mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}$ let $\mathrm{Perm}_{V,W} \colon V \otimes W \to W \otimes V$ be the permutation $\mathrm{Perm}_{V,W}(v \otimes w) = w \otimes v$ and define $\beta^R_{V,W} = \mathrm{Perm}_{V,W} \circ R$. We scale $\beta^R_{V,W}$ as follows:

$$\tilde{\beta}^R_{V,W} = q^{-\frac{kl}{N}} \beta^R_{V,W}$$

whenever V and W are exterior or symmetric power $\mathbf{U}_q(\mathfrak{gl}_N)$-modules of exponent k and l respectively. This induces a scaling $\tilde{\beta}^R_{V,W}$ of $\beta^R_{V,W}$ for all $\mathbf{U}_q(\mathfrak{gl}_N)$-modules $V, W \in \mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}$. Then $(\mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}, \tilde{\beta}^R_{\cdot,\cdot})$ is a braided monoidal category.

The goal of this subsection is to finally prove our main theorems. To this end, we extend (21) to a diagram

$$\begin{tikzcd}
\dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n}) \arrow[r, "\Phi^{m|n}_{\mathrm{su}}"] \arrow[dr, "\Upsilon^{m|n}_{\mathrm{su}}"'] & \mathfrak{gl}_N\text{-}\mathbf{Mod}^{\mathrm{sort}}_{\mathrm{es}} \arrow[r, hook, "\iota_{\mathrm{alg}}"] & \mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}} \\
 & N\text{-}\mathbf{Web}^{\mathrm{sort}}_{\mathrm{gr}} \arrow[u, "\Gamma^{\mathrm{sort}}_N"'] \arrow[r, hook, "\iota_{\mathrm{dia}}"'] & N\text{-}\mathbf{Web}_{\mathrm{gr}} \arrow[u, "\Gamma_N"']
\end{tikzcd} \tag{23}$$

where ι_{alg} and ι_{dia} are the evident inclusions of full subcategories. We will define the functor Γ_N such that the diagram (23) commutes.

Definition 3.17. (**The diagrammatic presentation functor** Γ_N) We define a functor $\Gamma_N \colon N\text{-}\mathbf{Web}_{\mathrm{gr}} \to \mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}$ as follows.

- On objects: Γ_N sends an object $\vec{k} \in X^L$ of N-$\mathbf{Web}_{\mathrm{gr}}$ to the tensor product of exterior and symmetric powers of $\mathbb{C}_q^N$ specified by the entries of $\vec{k}$: green and red integers encode exterior and symmetric powers respectively, and a black entry 1 corresponds to $\mathbb{C}_q^N$ itself.
- On morphisms: For an object $\vec{k} \in X^L$ let $w(\vec{k}) \in S_L$ be a shortest length permutation that sorts green integers in $\vec{k}$ to the left of red integers. We define Γ_N on an arbitrary web $u \in \mathrm{Hom}_{N\text{-}\mathbf{Web}_{\mathrm{gr}}}(\vec{k}, \vec{l})$ by pre-composing and post-composing with elementary crossings and the universal R-matrix intertwiners:
$$\Gamma_N(u) = \tilde{\beta}^R_{w(\vec{l})^{-1}} \circ \Gamma_N^{\mathrm{sort}}(\beta^\bullet_{w(\vec{l})} \circ u \circ \beta^\bullet_{w(\vec{k})^{-1}}) \circ \tilde{\beta}^R_{w(\vec{k})}.$$

Clearly, Γ_N restricts to Γ_N^{sort}. ◇

Lemma 3.18. $\Gamma_N \colon N$-$\mathbf{Web}_{\mathrm{gr}} \to \mathfrak{gl}_N$-$\mathbf{Mod}_{\mathrm{es}}$ is a monoidal functor making (23) commutative.

Proof. By Lemma 3.12 and the fact that $\beta^\bullet_{\cdot,\cdot}$ and $\tilde{\beta}^R_{\cdot,\cdot}$ are braidings (see Proposition 2.22 and Remark 3.16), we see that Γ_N is well-defined. That Γ_N is monoidal and makes (23) commutative is clear from the construction. ☐

Proposition 3.19. The functor $\Gamma_N \colon (N\text{-}\mathbf{Web}_{\mathrm{gr}}, \beta^\bullet_{\cdot,\cdot}) \to (\mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}, \tilde{\beta}^R_{\cdot,\cdot})$ is a functor of braided monoidal categories.

Proof. By Lemma 3.18, it remains to verify
$$\Gamma_N(\beta^\bullet_{\vec{k}\otimes\vec{l}}) = \tilde{\beta}^R_{\Gamma_N(\vec{k}),\Gamma_N(\vec{l})}, \quad \text{for all objects } \vec{k}, \vec{l} \text{ of } N\text{-}\mathbf{Web}_{\mathrm{gr}}.$$

The green-red symmetry and the fact that the mixed crossings are defined via the monochromatic crossings, together with Corollary 2.24, reduce this problem to the situation studied in [4, Theorem 6.2.1 and Lemma 6.2.2]. It remains to show
$$\Gamma_N(\beta^g_{1,1}) = \Gamma_N(\beta^r_{1,1}) = \Gamma_N^{\mathrm{sort}}(\beta^g_{1,1}) = \Gamma_N^{\mathrm{sort}}(\beta^r_{1,1}) = \tilde{\beta}^R_{\mathbb{C}_q^N,\mathbb{C}_q^N}.$$

This follows since $\Gamma_N^{\mathrm{sort}}(\beta^g_{1,1}) = \Gamma_N^{\mathrm{sort}}(\beta^r_{1,1})$ acts on $\mathbb{C}_q^N \otimes \mathbb{C}_q^N \cong \textstyle\bigwedge_q^2(\mathbb{C}_q^N) \oplus \mathrm{Sym}_q^2(\mathbb{C}_q^N)$ as q on the first factor and as $-q^{-1}$ on the second (see Example 2.18). ☐

Theorem 3.20. (**The diagrammatic presentations**) The functor
$$\Gamma_N \colon (\mathbf{Mat}(N\text{-}\mathbf{Web}_{\mathrm{gr}}), \beta^\bullet_{\cdot,\cdot}) \to (\mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}, \tilde{\beta}^R_{\cdot,\cdot})$$
is an equivalence of braided monoidal categories.

Proof. By Proposition 3.19, Γ_N extends to a braided monoidal functor on the additive closure and it remains to show that Γ_N is essentially surjective, full and faithful.

Essentially surjective: this follows directly from the definitions. See also Remark 3.14.

Full and faithful: as before, it suffices to verify this on morphisms between objects of the form $\vec{k} \in X^L$. Consider the commuting diagram

$$\begin{tikzcd}
\mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}^{\mathrm{sort}} & \mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}} \arrow[l, "\omega_R"'] \\
N\text{-}\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}} \arrow[u, "\Gamma_N^{\mathrm{sort}}"] & N\text{-}\mathbf{Web}_{\mathrm{gr}} \arrow[l, "\omega_\bullet"] \arrow[u, "\Gamma_N"']
\end{tikzcd},$$

where ω_R and $\omega_\bullet$ are the functors that order $f \in \mathrm{Hom}_{\mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}}(\Gamma_N(\vec{k}),\Gamma_N(\vec{l}))$ and webs $u \in \mathrm{Hom}_{N\text{-}\mathbf{Web}_{\mathrm{gr}}}(\vec{k},\vec{l})$ by using the R-matrix braiding $\tilde{\beta}^{R}_{\cdot,\cdot}$ and the braiding $\beta^{\bullet}_{\cdot,\cdot}$ respectively, via a permutation of shortest length. Since sorting is invertible we get:

$$\begin{aligned}\dim(\mathrm{Hom}_{\mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}}(\Gamma_N(\vec{k}),\Gamma_N(\vec{l}))) &= \dim(\mathrm{Hom}_{\mathfrak{gl}_N\text{-}\mathbf{Mod}^{\mathrm{sort}}_{\mathrm{es}}}(\Gamma^{\mathrm{sort}}_N(\omega_\bullet(\vec{k})),\Gamma^{\mathrm{sort}}_N(\omega_\bullet(\vec{l}))))\\ &= \dim(\mathrm{Hom}_{N\text{-}\mathbf{Web}^{\mathrm{sort}}_{\mathrm{gr}}}(\omega_\bullet(\vec{k}),\omega_\bullet(\vec{l})))\\ &= \dim(\mathrm{Hom}_{N\text{-}\mathbf{Web}_{\mathrm{gr}}}(\vec{k},\vec{l})),\end{aligned}$$

where the second equality follows from Proposition 3.15. ☐

Remark 3.21. For now we restrict ourselves to working with webs with only upward oriented edges. Downward oriented edges, as for example in [4], can used to represent the duals of the $\mathbf{U}_q(\mathfrak{gl}_N)$-modules $\textstyle\bigwedge^k_q\mathbb{C}^N_q$ and $\mathrm{Sym}^l_q\mathbb{C}^N_q$.With respect to such an enriched web calculus, the statement of Theorem 3.20 extends to an equivalence of pivotal categories, see [21, Section 6] and Remark 5.12.

Let $\check{\mathbf{H}}$ denotes the monoidal, $\mathbb{C}_q$-linear category obtained from the collection $H_\infty(q)$ of Iwahori-Hecke algebras as follows. The objects e, e' of $\check{\mathbf{H}}$ are tensor products of Iwahori-Hecke algebra idempotents corresponding to $e_{\mathrm{col}}(\lambda)$ and $e_{\mathrm{row}}(\lambda)$ (as in Definition 2.26) under the isomorphism in Lemma 2.25. The morphism spaces are given by $\mathrm{Hom}_{\check{\mathbf{H}}}(e,e') = e'H_\infty(q)e$. The category $\check{\mathbf{H}}$ is braided with braiding $\tilde{\beta}^{H}_{\cdot,\cdot}$ induced from $H_\infty(q)$.

Theorem 3.22. (**The diagrammatic presentation**) For large N the functors Γ_N stabilize to a functor

$$\Gamma_\infty\colon (\mathbf{Mat}(\infty\text{-}\mathbf{Web}_{\mathrm{gr}}),\beta^{\bullet}_{\cdot,\cdot}) \to (\mathbf{Mat}(\check{\mathbf{H}}),\tilde{\beta}^{H}_{\cdot,\cdot}),$$

which is an equivalence braided monoidal categories.

Proof. By Schur-Weyl duality (15) and by the construction of the categories N-$\mathbf{Web}_{\mathrm{gr}}$ as quotients of ∞-$\mathbf{Web}_{\mathrm{gr}}$, we have quotient functors π^N_∞ and π^N for $N \in \mathbb{Z}_{\geq 0}$ such that

$$\begin{tikzcd} \mathbf{Mat}(\check{\mathbf{H}}) \arrow[r, "\pi^N"] & \mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}} \\ \mathbf{Mat}(\infty\text{-}\mathbf{Web}_{\mathrm{gr}}) \arrow[u, "\Gamma_\infty"] \arrow[r, "\pi^N_\infty"'] & \mathbf{Mat}(N\text{-}\mathbf{Web}_{\mathrm{gr}}) \arrow[u, "\Gamma_N"'] \end{tikzcd} \tag{24}$$

commutes. Here the functor Γ_∞ is an idempotented version of the inverse of the isomorphism $\Phi^{\infty}_{q\mathrm{SW}}$ from Lemma 2.25.

Fix two objects $\vec{k} \in X^L$ and $\vec{l} \in X^L$ of ∞-$\mathbf{Web}_{\mathrm{gr}}$ and suppose that N is greater than the sum of the integer-values of the entries of $\vec{k}$ (ignoring their colors). Then, by (15), Theorem 3.20, the commutativity of (24) and the fullness of π^N_∞, we have

$$\begin{aligned}\dim(\mathrm{Hom}_{\check{\mathbf{H}}}(\Gamma_\infty(\vec{k}),\Gamma_\infty(\vec{l}))) &= \dim(\mathrm{Hom}_{\mathfrak{gl}_N\text{-}\mathbf{Mod}_{\mathrm{es}}}(\pi^N(\Gamma_\infty(\vec{k})),\pi^N(\Gamma_\infty(\vec{l}))))\\ &= \dim(\mathrm{Hom}_{N\text{-}\mathbf{Web}_{\mathrm{gr}}}(\pi^N_\infty(\vec{k}),\pi^N_\infty(\vec{l})))\\ &= \dim(\mathrm{Hom}_{\infty\text{-}\mathbf{Web}_{\mathrm{gr}}}(\vec{k},\vec{l})).\end{aligned}$$

Γ_∞ is clearly essentially surjective and a braided monoidal functor and the theorem follows. ☐

4. Applications

In this section we write $\mathcal{L}_D$ for *diagrams of framed, oriented links* $\mathcal{L}$, b_D^K for *diagrams of braids in K-strands* and $\overline{b}_D^K$ for *closures* of such braid diagrams. Further, we consider labelings of the connected components of $\mathcal{L}$ and of braids by Young diagrams λ^i. If $\mathcal{L}$ is a d-component link, then we write $\mathcal{L}(\vec{\lambda})$ for its labeling by a vector of Young diagrams $\vec{\lambda} = (\lambda^1, \dots, \lambda^d)$, and use an analogous notation for labeled link and braid diagrams. If not mentioned otherwise, then all appearing links and related concepts are assumed to be framed and oriented from now on.

Let $\mathcal{L}_D(\vec{\lambda}) = \overline{b}_D^K(\vec{\lambda})$ be a diagram of a framed, oriented, labeled link given as a braid closure. The following process associates to $b_D^K(\vec{\lambda})$ an element $p_{K'}(\tilde{b}_D^{K'})$ of $H_{K'}(q) \cong \mathrm{End}_{\infty\text{-}\mathbf{Web}_{\mathrm{gr}}}(\vec{K}')$:



where we write p_{K_i} for the Hecke algebra representation of the braid group on K_i strands. The first step replaces strands labeled by a Young diagram λ^i with K_i nodes in the braid diagram b_D^K by K_i parallel strands. This results in a new braid $\tilde{b}_D^{K'}$ where K' indicates the number of strands. In the second step this cabled braid is interpreted as an element of the Hecke algebra, or equivalently, as a web in ∞-$\mathbf{Web}_{\mathrm{gr}}$, with an idempotent $e_q(\lambda^i)$ placed on the cable of each previously λ^i labeled strand.

4.1. The colored HOMFLY-PT polynomial via the category ∞-$\mathbf{Web}_{\mathrm{gr}}$.

In this subsection we work over the ground field $\mathbb{C}_{a,q} = \mathbb{C}_q(a)$. We will use the $\mathbb{C}_{a,q}$-valued *Jones-Ocneanu trace* $\mathrm{tr}(\cdot)$ on the collection of Iwahori-Hecke algebras $H_\infty(q) = \bigoplus_{K \in \mathbb{Z}_{\geq 0}} H_K(q)$. The definition of the trace $\mathrm{tr}(\cdot)$ can be found in [13, Section 5] (which can be easily adopted to our notation). We will use it in the form of the following lemma.

Lemma 4.1. Given a web $u \in \mathrm{End}_{\infty\text{-}\mathbf{Web}_{\mathrm{gr}}}(\vec{K})$. Then

$\mathrm{tr}(u) =$



$\in \mathbb{C}_{a,q}$,

where the diagram can be collapsed by using the relations in ∞-$\mathbf{Web}_{\mathrm{gr}}$ and additionally

(25) 

$= \frac{a-a^{-1}}{q-q^{-1}}$, 

$= \frac{aq^{-1}-a^{-1}q}{q-q^{-1}}$ 

Proof. By Proposition 2.15 and Corollary 2.13: any given web $u \in \mathrm{End}_{\infty\text{-}\mathbf{Web}_{\mathrm{gr}}}(\vec{K})$ can be expressed using black or green edges with labels at most 2. Using Lemma 2.25 and additionally [22, Subsection 4.2], where Rasmussen's singular crossings correspond to green dumbbells with label 2, provides the statement. Note that Rasmussen's relations II and III are already part of our diagrammatic calculus. □

Definition 4.2. (**The colored HOMFLY-PT polynomial**) Let $\mathcal{L}_D(\vec{\lambda}) = \overline{b}_D^K(\vec{\lambda})$ be a diagram of a framed, oriented, labeled link $\mathcal{L}(\vec{\lambda})$ given as a braid closure.

The *colored HOMFLY-PT polynomial* of $\mathcal{L}(\vec{\lambda})$, denoted by $\mathcal{P}^{a,q}(\mathcal{L}(\vec{\lambda}))$, is defined via

$$\mathcal{P}^{a,q}(\mathcal{L}(\vec{\lambda})) = \mathrm{tr}(p_{K'}(\tilde{b}_D^{K'})e_q(\vec{\lambda})) \in \mathbb{C}_{a,q},$$

where $e_q(\vec{\lambda})$ is a tensor product of $e_q(\lambda^i)$'s, as described above. ◇

This polynomial is independent of all choices involved and an invariant of framed, oriented, colored links. Up to different conventions, this is shown for example in [29, Corollary 4.5].

Remark 4.3. In fact, Definition 4.2 gives the framing dependent, unnormalized version of the colored HOMFLY-PT polynomial. As usual, the polynomial can be normalized by fixing the value of the unknot to be 1 (instead of $\frac{a-a^{-1}}{q-q^{-1}}$ as in our convention) and one can get rid of the framing dependence by scaling with a factor coming from Reidemeister 1 moves, see for example [13, Definition 6.1]. We suppress these distinctions in the following.

Note that Lemma 4.1 provides a method to calculate the colored HOMFLY-PT polynomials $\mathcal{P}^{a,q}(\cdot)$ using the web category ∞-$\mathbf{Web}_{\mathrm{gr}}$.

Proposition 4.4. (**The colored HOMFLY-PT symmetry**) We have

$$\mathcal{P}^{a,q}(\mathcal{L}(\vec{\lambda})) \;=\; (-1)^{co}\,\mathcal{P}^{a,q^{-1}}(\mathcal{L}(\vec{\lambda}^{\,\mathrm{T}})),$$

where $\vec{\lambda}^{\,\mathrm{T}} = ((\lambda^1)^{\mathrm{T}}, \dots, (\lambda^d)^{\mathrm{T}})$. Moreover, $co \in \mathbb{Z}$ is a certain constant depending only on the framed, oriented link $\mathcal{L}$ and its coloring.

This symmetry is not new: it can be deduced from [18, Section 9] and has been studied in [17]. In our framework it follows directly from the green-red symmetry in ∞-$\mathbf{Web}_{\mathrm{gr}}$.

Proof. We only give a proof for the case of knots $\mathcal{K}$. The proof for links is analogous, but the notation is more involved. We denote by I_{gr} the involution on ∞-$\mathbf{Web}_{\mathrm{gr}}$ given by the green-red symmetry, and by I_q the involution on $\mathbb{C}_{a,q}$ which inverts the variable q.

Claim: for $u \in \mathrm{End}_{\infty\text{-}\mathbf{Web}_{\mathrm{gr}}}(\vec{K})$ we have

$$\mathrm{tr}(u) = (-1)^K I_q(\mathrm{tr}(I_{\mathrm{gr}}(u))). \tag{26}$$

It suffices to prove this in the case where u is a primitive web (a morphism that consists of a single web with coefficient 1). In Lemma 4.1 we have met evaluation relations for monochromatic green webs of edge label at most 2, but clearly analogous relations can be derived for red and mixed webs. In fact, all necessary evaluation relations are invariant under I_{gr} and I_q, except the two relations in (25). The circle relation is I_{gr} invariant, but acquires a sign under I_q. The following computation shows that the green and red bubble relations also respect (26):


1
1
2
1
$\overset{(10)}{=}$
$[2]$
1
1
1
1
$-$
2
1
$=$
$\dfrac{aq-a^{-1}q^{-1}}{q-q^{-1}}$
1
1
$\xleftrightarrow{q\leftrightarrow q^{-1}}$
$-$
1
1
2
1

We note that in the computation of $\mathrm{tr}(u)$ via Lemma 4.1 strands can only be removed by circle moves and bubble moves. Both of these acquire a sign under I_q, which causes the factor $(-1)^K$ in (26). This proves the claim.

Let b_D^K be a braid diagram that closes to a diagram of $\mathcal{K}$ and suppose that $\mathcal{K}$ is labeled by a Young diagram λ of with L nodes. Let $\tilde{b}_D^{KL}$ be the L-fold cable of the braid diagram b_D^K.

Now we have

$$\begin{aligned}\mathcal{P}^{a,q}(\mathcal{K}(\lambda)) = \mathrm{tr}\left(p_{KL}\left(\tilde{b}_D^{KL}\right)e_q(\lambda)^{\otimes K}\right) &= (-1)^{KL} I_q\left(\mathrm{tr}\left(I_{\mathrm{gr}}\left(p_{KL}\left(\tilde{b}_D^{KL}\right)e_q(\lambda)^{\otimes K}\right)\right)\right)\\ &= (-1)^{(\mathrm{cr}+1)KL} I_q\left(\mathrm{tr}\left(p_{KL}\left(\tilde{b}_D^{KL}\right)e_q(\lambda^{\mathrm{T}})^{\otimes K}\right)\right)\\ &= (-1)^{(\mathrm{cr}+1)KL}\,\mathcal{P}^{a,q^{-1}}(\mathcal{K}(\lambda^{\mathrm{T}})),\end{aligned}$$

where cr is the number of crossings of b_D^K. Here we have used (26), Lemma 2.28 (note that conjugation by a braid does not matter inside the trace) and the fact that I_{gr} acts as -1 on black crossings, see Example 2.18 □

4.2. **The colored $\mathfrak{sl}_N$-link polynomials via the categories N-Web$_{\mathrm{gr}}$.** Recall that the *colored Reshetikhin-Turaev $\mathfrak{sl}_N$-link polynomial* $\mathcal{RT}^{q^N,q}(\mathcal{L}(\vec{\lambda}))$ are determined by the corresponding colored HOMFLY-PT polynomials $\mathcal{P}^{a,q}(\mathcal{L}(\vec{\lambda}))$ by specializing $a = q^N$. Alternatively, they can be computed directly inside the categories N-**Web**$_{\mathrm{gr}}$ from a framed, oriented, labeled link diagram as follows:

- First we replace all λ-labeled strands in the link diagram by cables equipped with the diagrammatic idempotent $e_q(\lambda)$, written in monochromatic green webs.
- The resulting diagram will contain downward oriented green edges of label k, which we replace by upward oriented green edges of label $N-k$. Simultaneously, caps and cups are replaced by splits and merges



- The result is a morphism in N-$\mathbf{Web}_{\mathrm{gr}}$ between objects consisting only of entries 0 and N_g. It follows from Theorem 3.20 that this Hom-space is one-dimensional. Thus, the framed, oriented, labeled link diagram determines a polynomial, which is the desired colored Reshetikhin-Turaev $\mathfrak{sl}_N$-link polynomial.

Recall from Remark 1.1 that this approach relies on the fact that $\mathfrak{sl}_N$-$\mathbf{Mod}_{\mathrm{es}}$ contains the duality isomorphisms $\textstyle\bigwedge_q^k\mathbb{C}_q^N \cong (\bigwedge_q^{N-k}\mathbb{C}_q^N)^*$, which is not the case for $\mathfrak{gl}_N$-$\mathbf{Mod}_{\mathrm{es}}$ or $\mathfrak{gl}_{N|M}$-$\mathbf{Mod}_{\mathrm{es}}$. In Remark 5.12 we sketch how to include duals in diagrammatic presentations of $\mathfrak{gl}_N$-$\mathbf{Mod}_{\mathrm{es}}$ and $\mathfrak{gl}_{N|M}$-$\mathbf{Mod}_{\mathrm{es}}$ and, thus, to compute the corresponding Reshetikhin-Turaev $\mathfrak{gl}_N$ or $\mathfrak{gl}_{m|n}$-link invariants.

5. GENERALIZATION TO WEBS FOR $\mathfrak{gl}_{N|M}$

We now give a diagrammatic presentation of $\mathfrak{gl}_{N|M}$-$\mathbf{Mod}_{\mathrm{es}}$, the braided monoidal category of $\mathbf{U}_q(\mathfrak{gl}_{N|M})$-modules tensor generated by the exterior powers $\textstyle\bigwedge_q^k\mathbb{C}_q^{N|M}$ and the symmetric powers $\mathrm{Sym}_q^l\mathbb{C}_q^{N|M}$ of the vector representation $\mathbb{C}_q^{N|M}$ of $\mathbf{U}_q(\mathfrak{gl}_{N|M})$. The diagrammatic presentation is given by the following quotient of ∞-$\mathbf{Web}_{\mathrm{gr}}$.

Definition 5.1. The category $N|M$-$\mathbf{Web}_{\mathrm{gr}}$ is the quotient category obtained from ∞-$\mathbf{Web}_{\mathrm{gr}}$ by imposing the *not-a-hook relation*, that is

$$e_q(\mathrm{box}_{N+1,M+1}) = 0.$$

Where $\mathrm{box}_{N+1,M+1}$ is the box-shaped Young diagram with $N+1$ rows and $M+1$ columns. ◇

Note that $N|M$-$\mathbf{Web}_{\mathrm{gr}}$ inherits the braiding $\beta^{\bullet}_{\cdot,\cdot}$ from ∞-$\mathbf{Web}_{\mathrm{gr}}$.

Example 5.2. If we take $M=0$, then $\mathrm{box}_{N+1,1}$ is a column Young diagram with $N+1$ nodes and the corresponding not-a-hook relation is just the exterior relation (11). In this case we have that $N|0$-$\mathbf{Web}_{\mathrm{gr}}$ is N-$\mathbf{Web}_{\mathrm{gr}}$ and $\mathfrak{gl}_{N|0}$-$\mathbf{Mod}_{\mathrm{es}}$ is isomorphic to $\mathfrak{gl}_N$-$\mathbf{Mod}_{\mathrm{es}}$. ■

Example 5.3. If we take $M=N=1$, then we have



It is easy to see that $e_q(\mathrm{box}_{2,2}) = 0$ is equivalent to the relations [26, (3.3.13a) and (3.3.13b)], [8, Subsection 3.6] and [21, Corollary 6.18] which are used to describe the "purely exterior" representation category $\mathfrak{gl}_{1|1}$-$\mathbf{Mod}_{\mathrm{e}}$. This category could be presented as monochromatic green subcategory of $1|1$-$\mathbf{Web}_{\mathrm{gr}}$, defined analogously as in Definition 2.7. ■

To prove that $N|M$-$\mathbf{Web}_{\mathrm{gr}}$ gives a diagrammatic presentation of $\mathfrak{gl}_{N|M}$-$\mathbf{Mod}_{\mathrm{es}}$, we use a version of super q-Howe duality between $\mathbf{U}_q(\mathfrak{gl}_{m|n})$ and $\mathbf{U}_q(\mathfrak{gl}_{N|M})$. For this, we say a dominant integral $\mathfrak{gl}_{m|n}$-weight λ is $(m|n,M|N)$-*supported* if it corresponds to a Young diagram which is simultaneously an $(m|n)$-hook as well as an $(M|N)$-hook[8].

---

[8] This is really intended to be $(M|N)$.

Theorem 5.4. (**Super q-Howe duality, super-super version**) We have the following.

(a) Let $K \in \mathbb{Z}_{\geq 0}$. The actions of $\mathbf{U}_q(\mathfrak{gl}_{m|n})$ and $\mathbf{U}_q(\mathfrak{gl}_{N|M})$ on ${\textstyle\bigwedge}_q^K(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^{N|M})$ commute and generate each others commutant.

(b) There exists an isomorphism

$$ {\textstyle\bigwedge}_q^{\bullet}(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^{N|M}) \cong ({\textstyle\bigwedge}_q^{\bullet}\mathbb{C}_q^{N|M})^{\otimes m} \otimes (\mathrm{Sym}_q^{\bullet}\mathbb{C}_q^{N|M})^{\otimes n} $$

of $\mathbf{U}_q(\mathfrak{gl}_{N|M})$-modules under which the $\vec{k}$-weight space of ${\textstyle\bigwedge}_q^{\bullet}(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^{N|M})$ (considered as a $\mathbf{U}_q(\mathfrak{gl}_{m|n})$-module) is identified with

$$ {\textstyle\bigwedge}_q^{\vec{k}_0}\mathbb{C}_q^{N|M} \otimes \mathrm{Sym}_q^{\vec{k}_1}\mathbb{C}_q^{N|M} = \\ {\textstyle\bigwedge}_q^{k_1}\mathbb{C}_q^{N|M} \otimes \cdots \otimes {\textstyle\bigwedge}_q^{k_m}\mathbb{C}_q^{N|M} \otimes \mathrm{Sym}_q^{k_{m+1}}\mathbb{C}_q^{N|M} \otimes \cdots \otimes \mathrm{Sym}_q^{k_{m+n}}\mathbb{C}_q^{N|M}. $$

Here $\vec{k} = (k_1, \dots, k_{m+n})$, $\vec{k}_0 = (k_1, \dots, k_m)$ and $\vec{k}_1 = (k_{m+1}, \dots, k_{m+n})$.

(c) As $\mathbf{U}_q(\mathfrak{gl}_{m|n}) \otimes \mathbf{U}_q(\mathfrak{gl}_{N|M})$-modules, we have a decomposition of the form

$$ {\textstyle\bigwedge}_q^K(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^{N|M}) \cong \bigoplus_{\lambda} L_{m|n}(\lambda) \otimes L_{N|M}(\lambda^{\mathrm{T}}), $$

where we sum over all $(m|n, M|N)$-supported $\mathfrak{gl}_{m|n}$-weights λ whose entries sum up to K. This induces a decomposition

$$ {\textstyle\bigwedge}_q^{\bullet}(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^{N|M}) \cong \bigoplus_{\lambda} L_{m|n}(\lambda) \otimes L_{N|M}(\lambda^{\mathrm{T}}), $$

where we sum over all $(m|n, M|N)$-supported $\mathfrak{gl}_{m|n}$-weights λ.

Proof. As before, (a) and (c) are proven in [21, Theorem 4.2] and only (b) remains to be verified. This works similarly as in the proof of Theorem 3.6 and is left to the reader. For a non-quantized version see [25, Proposition 2.2]. □

In the statement of this theorem, ${\textstyle\bigwedge}_q^k\mathbb{C}_q^{N|M}$, $\mathrm{Sym}_q^l\mathbb{C}_q^{N|M}$ and ${\textstyle\bigwedge}_q^K(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^{N|M})$ are defined similarly as in Subsection 3.1, see also [21, Section 3]. As before we then get:

Corollary 5.5. There exists a full functor $\Phi_{\mathrm{su}}^{m|n} \colon \dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n}) \to \mathfrak{gl}_{N|M}\text{-}\mathbf{Mod}_{\mathrm{es}}$, which we again call the *super q-Howe functor*, given on objects and morphisms by

$$ \vec{k} \overset{\Phi_{\mathrm{su}}^{m|n}}{\longmapsto} {\textstyle\bigwedge}_q^{\vec{k}_0}\mathbb{C}_q^{N|M} \otimes \mathrm{Sym}_q^{\vec{k}_1}\mathbb{C}_q^{N|M} \quad , \quad 1_{\vec{l}}x1_{\vec{k}} \overset{\Phi_{\mathrm{su}}^{m|n}}{\longmapsto} f_{\vec{k}}^{\vec{l}}(x). $$

Everything else is sent to zero. □

In what follows, we denote by $\dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n})^{\geq 0}$ the quotient of $\dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n})$ obtained by killing all $\mathfrak{gl}_{m|n}$-weights with negative entries.

Corollary 5.6. The super q-Howe functor $\Phi_{\mathrm{su}}^{m|n}$ from Corollary 5.5 induces an algebra epimorphism (denoted by the same symbol) as in the diagram below. Under Artin-Wedderburn decompositions, it corresponds to an algebra epimorphism π, which acts on the summand

$\mathrm{End}_{\mathbb{C}_q}(L_{m|n}(\lambda))$ either as an isomorphism or as zero, depending on whether the Young diagram λ is $(m|n, M|N)$-supported or not.

$$\begin{tikzcd}
\dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n})^{\geq 0} \arrow[r, "\cong"] \arrow[d, two heads, "\Phi_{\mathrm{su}}^{m|n}"'] & \bigoplus\limits_{\substack{(m|n)\text{-}\\ \text{hooks } \lambda}} \mathrm{End}_{\mathbb{C}_q}(L_{m|n}(\lambda)) \arrow[d, two heads, "\pi"] \\
\mathrm{End}_{\mathbf{U}_q(\mathfrak{gl}_{N|M})}(\textstyle\bigwedge_q^{\bullet}(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^{N|M})) \arrow[r, "\cong"'] & \bigoplus\limits_{\substack{(m|n,M|N)\text{-}\\ \text{supported } \lambda}} \mathrm{End}_{\mathbb{C}_q}(L_{m|n}(\lambda))
\end{tikzcd}$$

Proof. First, note that by Theorem 3.22, $\dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n})^{\geq 0}$ is isomorphic to $\check{\mathbf{H}}_{m+n}^{\mathrm{sort}}$, the sorted version of $\check{\mathbf{H}}$ with exactly m exterior strands and n symmetric strands. The Artin-Wedderburn decomposition in the top row of the diagram is then given in [19, Theorem 5.1]. The bottom Artin-Wedderburn decomposition follows directly from part (c) of Theorem 5.4. ☐

Remark 5.7. We obtain from Corollary 5.6 an alternative proof of the presentation of the q-Schur superalgebra $S_q(N|M, K) \cong \mathrm{End}_{H_K(q)}((\mathbb{C}_q^{N|M})^{\otimes K})$ from [6, Theorem 3.13.1].

Lemma 5.8. Under the correspondence

$$\check{\mathbf{H}}_{m+n}^{\mathrm{sort}} \xleftrightarrow{\cong} \dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n})^{\geq 0} \xleftrightarrow{\cong} \bigoplus_{\substack{(m|n)\text{-}\\ \text{hooks } \lambda}} \mathrm{End}_{\mathbb{C}_q}(L_{m|n}(\lambda))$$

the kernel of the super q-Howe functor $\Phi_{\mathrm{su}}^{m|n}$ from Corollary 5.5 is given by the tensor ideal I_{box} in $\check{\mathbf{H}}_{m+n}^{\mathrm{sort}}$ generated by the primitive idempotent $e_q(\mathrm{box}_{N+1,M+1})$.

Proof. From the right isomorphism of algebras we know that the kernel of $\Phi_{\mathrm{su}}^{m|n}$ is generated by all $e_q(\lambda^{\mathrm{T}})$ where λ is an $(m|n)$-hook, but not an $(M|N)$-hook. Note that every such λ corresponds to a simple $\mathbf{U}_q(\mathfrak{gl}_{N|M})$-module which appears in a tensor product $L_{N|M}((\mathrm{box}_{N+1,M+1})^{\mathrm{T}}) \otimes (\mathbb{C}_q^{N|M})^{\otimes K}$ for some $K \in \mathbb{Z}_{\geq 0}$. Accordingly, $e_q(\lambda^T)$ is contained in the ideal I_{box}. ☐

Proposition 5.9. There is an equivalence of categories

$$\mathbf{Mat}(N|M\text{-}\mathbf{Web}_{\mathrm{gr}}^{\mathrm{sort}}) \cong \mathfrak{gl}_{N|M}\text{-}\mathbf{Mod}_{\mathrm{es}}^{\mathrm{sort}}.$$

Proof. Lemma 5.8 shows that the sorted web category $N|M\text{-}\mathbf{Web}_{m+n}^{\mathrm{sort}}$, in which webs have m green and n red boundary points both on the bottom and on the top, is equivalent to $\mathrm{End}_{\mathbf{U}_q(\mathfrak{gl}_{N|M})}(\bigwedge_q^{\bullet}(\mathbb{C}_q^{m|n} \otimes \mathbb{C}_q^{N|M}))$, considered as a category. Via the $\dot{\mathbf{U}}_q(\mathfrak{gl}_{m|n})$-weight space decomposition in part (b) of Theorem 5.4, $N|M\text{-}\mathbf{Web}_{m+n}^{\mathrm{sort}}$ gives a presentation of the morphism spaces in $\mathfrak{gl}_{N|M}\text{-}\mathbf{Mod}_{\mathrm{es}}^{\mathrm{sort}}$ between objects of the form

$$\textstyle\bigwedge_q^{k_1}\mathbb{C}_q^{N|M} \otimes \cdots \otimes \bigwedge_q^{k_m}\mathbb{C}_q^{N|M} \otimes \mathrm{Sym}_q^{k_{m+1}}\mathbb{C}_q^{N|M} \otimes \cdots \otimes \mathrm{Sym}_q^{k_{m+n}}\mathbb{C}_q^{N|M}.$$

Any object in $\mathfrak{gl}_{N|M}\text{-}\mathbf{Mod}_{\mathrm{es}}^{\mathrm{sort}}$ is a formal sum of such objects, for suitable $m, n \in \mathbb{Z}_{\geq 0}$, and the conclusion follows. ☐

Remark 5.10. Recall that the category $\mathfrak{gl}_{N|M}$-$\mathbf{Mod}_{\mathrm{es}}$ is a braided monoidal category, where the braiding $\beta^R_{\cdot,\cdot}$ is given by the *universal R-matrix for* $\mathfrak{gl}_{N|M}$, see [30]. As before, we use a rescaled braiding $\tilde{\beta}^R_{\cdot,\cdot}$, where we follow the conventions from [21, (3.11)] except that we substitute q by $-q$ in their formulas. In particular, $\tilde{\beta}^R_{\mathbb{C}_q^{N|M},\mathbb{C}_q^{N|M}}$ acts as q on $\bigwedge_q^2\mathbb{C}_q^{N|M}$ and as $-q^{-1}$ on $\mathrm{Sym}_q^2\mathbb{C}_q^{N|M}$.

Theorem 5.11. (**The diagrammatic presentation**) There is an equivalence of braided monoidal categories

$$(\mathbf{Mat}(N|M\text{-}\mathbf{Web}_{\mathrm{gr}}), \beta^{\bullet}_{\cdot,\cdot}) \cong (\mathfrak{gl}_{N|M}\text{-}\mathbf{Mod}_{\mathrm{es}}, \beta^R_{\cdot,\cdot}).$$

Proof. The equivalence from Proposition 5.9 can be extended to a monoidal functor between $\mathbf{Mat}(N|M\text{-}\mathbf{Web}_{\mathrm{gr}})$ and $\mathfrak{gl}_{N|M}$-$\mathbf{Mod}_{\mathrm{es}}$ as in Definition 3.17. We can also copy the proof of Proposition 3.19, where we use Remark 5.10 to prove that this functor respects the braiding. Equivalence via this functor follows then as in Theorem 3.20. □

Remark 5.12. In [21, Section 6] the authors show how to extend a diagrammatic presentation of $\mathfrak{gl}_{N|M}$-$\mathbf{Mod}_{\mathrm{e}}$ to diagrammatically encode the full subcategory of $\mathbf{U}_q(\mathfrak{gl}_{N|M})$-modules tensor generated by exterior powers and their duals. Graphically, this involves the introduction of additional objects corresponding to the duals of exterior powers, downward oriented edges (to represent identity morphisms on duals) and cap and cup webs (which represent co-evaluation and evaluation morphisms). Additional web relations including analogues of (25) are introduced to encode basic relationships between exterior powers and their duals. The extension of the diagrammatic presentation to include duals is then tautological and [21, Theorem 6.5] and [21, Proposition 6.16] show that the extended presentation functor is fully faithful.

They further show in [21, Proposition 6.15] that their graphical calculus allows the computation of the Reshetikhin-Turaev $\mathfrak{gl}_{N|M}$-tangle invariants for tangles labeled with exterior powers of the vector representation.

The same *spiderization* strategy – with minimal changes in proofs – gives an extension of our diagrammatic presentation $N|M$-$\mathbf{Web}_{\mathrm{gr}}$ of $\mathfrak{gl}_{N|M}$-$\mathbf{Mod}_{\mathrm{es}}$ to one for the full subcategory of $\mathbf{U}_q(\mathfrak{gl}_{N|M})$-modules tensor generated by exterior and symmetric powers and their duals. This spiderized green-red web category directly allows the computation of Reshetikhin-Turaev $\mathfrak{gl}_{N|M}$-tangle invariants for tangles labeled with exterior as well as symmetric powers of the vector representation. The cabling strategy from Section 4 can then be used to compute these invariants with respect to arbitrary irreducible representations.

We finish with the following corollary, which is a direct consequence of the discussion in this section and Proposition 4.4. More specifically, it is based on the facts that $N|M$-$\mathbf{Web}_{\mathrm{gr}}$ is defined as quotient of ∞-$\mathbf{Web}_{\mathrm{gr}}$ and that the spiderization in [21, Section 6] respects the specialization $a = q^{N-M}$ of the relations (25), which are sufficient to compute colored HOMFLY-PT polynomials of braid closures.

Corollary 5.13. We have:

(1) The Reshetikhin-Turaev $\mathfrak{gl}_{N|M}$-tangle invariant of a labeled tangle depends only on $N - M$. In the case of a labeled link, they agree with the specialization $a = q^{N-M}$ of the corresponding colored HOMFLY-PT polynomial.

(2) The green-red symmetry on ∞-$\mathbf{Web}_{\mathrm{gr}}$ descends to a symmetry between $N|M$-$\mathbf{Web}_{\mathrm{gr}}$ and $M|N$-$\mathbf{Web}_{\mathrm{gr}}$. Correspondingly, there is a symmetry between the representation categories of $\mathbf{U}_q(\mathfrak{gl}_{N|M})$ and $\mathbf{U}_q(\mathfrak{gl}_{M|N})$ that transposes Young diagrams indexing irreducibles.

(3) The symmetry of HOMFLY-PT polynomials described in Proposition 4.4 is a stabilized version of the symmetry between colored Reshetikhin-Turaev $\mathfrak{gl}_{N|M}$-link invariants and $\mathfrak{gl}_{M|N}$-link invariants which transposes Young diagrams and inverts q. $\square$

This confirms decategorified analogues of predictions about relationships between colored HOMFLY-PT homology and conjectural colored $\mathfrak{gl}_{N|M}$-link homologies, see [9].

REFERENCES

[1] A.K. Aiston and H.R. Morton, Idempotents of Hecke algebras of type A, J. Knot Theory Ramifications 7-4 (1998), 463-487, online available arXiv:q-alg/9702017.

[2] A. Beilinson, G. Lusztig and R. MacPherson, A geometric setting for the quantum deformation of $\mathfrak{gl}_n$, Duke Math. J. 61-2 (1990), 655-677.

[3] A. Berenstein and S. Zwicknagl, Braided symmetric and exterior algebras, Trans. Amer. Math. Soc. 360-7 (2008), 3429-3472, online available arXiv:math/0504155.

[4] S. Cautis, J. Kamnitzer and S. Morrison, Webs and quantum skew Howe duality, Math. Ann. 360-1-2 (2014), 351-390, online available arXiv:1210.6437.

[5] S.-J. Cheng and W. Wang, Howe duality for Lie superalgebras, Compositio Math. 128-1 (2001), 55-94, online available arXiv:math/0008093.

[6] H. El Turkey and J. Kujawa, Presenting Schur superalgebras, Pacific J. Math. 262-2 (2013), 285-316, online available arXiv:1209.6327.

[7] J. Grant, A Categorification of the skew Howe action on a representation category of $\mathbf{U}_q(\mathfrak{gl}_{m|n})$, in preparation.

[8] J. Grant, A generators and relations description of a representation category of $\mathbf{U}_q(\mathfrak{gl}(1|1))$, online available arXiv:1406.5399.

[9] E. Gorsky, S. Gukov and M. Stošić, Quadruply-graded colored homology of knots, online available arXiv:1304.3481.

[10] S. Gukov and M. Stošić, Homological algebra of knots and BPS states, Geom. Topol. Monogr. 18 (2012), 309-367, online available arXiv:1112.0030.

[11] A. Gyoja, A q-analogue of Young symmetrizer, Osaka J. Math. 23-4 (1986), 841-852.

[12] R. Howe, *Perspectives on invariant theory: Schur duality, multiplicity-free actions and beyond*, The Schur lectures, Israel Math. Conf. Proc. 8 (1992), 1-182.

[13] V.F.R. Jones, Hecke algebra representations of braid groups and link polynomials, Ann. of Math. (2) 126-2 (1987), 335-388.

[14] L.H. Kauffman and S.L. Lins, *Temperley-Lieb recoupling theory and invariants of* 3*-manifolds*, Annals of Mathematics Studies 134, Princeton University Press (1994).

[15] G. Kuperberg, Spiders for rank 2 Lie algebras, Comm. Math. Phys. 180-1 (1996), 109-151, online available arXiv:q-alg/9712003.

[16] J.-H. Kwon, Crystal bases of q-deformed Kac modules over the quantum superalgebras $\mathbf{U}_q(\mathfrak{gl}_{m|n})$, Int. Math. Res. Not. IMRN 2014-2, 512-550, online available arXiv:1203.5590.

[17] K. Liu and P. Peng, Proof of the Labastida-Marino-Ooguri-Vafa Conjecture, J. Differential Geom. 85-3 (2010), 479-525, online available arXiv:0704.1526.

[18] S.G. Lukac, Homfly skeins and the Hopf link, Ph.D. thesis, University of Liverpool (2001), online available http://www.liv.ac.uk/ su14/papers/lukacthesis.pdf.

[19] H. Mitsuhashi, Schur-Weyl reciprocity between the quantum superalgebra and the Iwahori-Hecke algebra, Algebr. Represent. Theory 9-3 (2006), 309-322, online available arXiv:math/0506156.

[20] H. Murakami, T. Ohtsuki and S. Yamada, HOMFLY polynomial via an invariant of colored plane graph, Enseign. Math. (2) 44-3-4 (1998), 325-360.

[21] H. Queffelec and A. Sartori, Mixed skew Howe duality and link invariants, online available arXiv:1504.01225.

[22] J. Rasmussen, Some differentials on Khovanov-Rozansky homology, to appear in Geom. Topol., online available arXiv:math/0607544.

[23] D.E.V. Rose and D. Tubbenhauer, Symmetric webs, Jones-Wenzl recursions and q-Howe duality, online available arXiv:1501.00915.

[24] G. Rumer, E. Teller and H. Weyl, Eine für die Valenztheorie geeignete Basis der binären Vektorinvarianten, Nachrichten von der Ges. der Wiss. Zu Göttingen, Math.-Phys. Klasse (1932), 498-504 (in German).

[25] A. Sartori, Categorification of tensor powers of the vector representation of $\mathbf{U}_q(\mathfrak{gl}(1|1))$, online available arXiv:1305.6162.

[26] A. Sartori, Categorification of tensor powers of the vector representation of $\mathbf{U}_q(\mathfrak{gl}(1|1))$, Ph.D. thesis, Universiät Bonn, online available http://hss.ulb.uni-bonn.de/2014/3635/3635.pdf.

[27] V. Turaev, *Quantum invariants of knots and* 3*-manifolds*, second revised edition, de Gruyter Studies in Mathematics (2010).

[28] Y. Wu and R.B. Zhang, Unitary highest weight representations of quantum general linear superalgebra, J. Algebra 321-11 (2009), 3568-3593.

[29] L. Xiao-Song and H. Zheng, On the Hecke algebras and the colored HOMFLY polynomial, Trans. Amer. Math. Soc. 362-1 (2010), 1-18, online available arXiv:math/0601267.

[30] H. Yamane, Quantized enveloping algebras associated with simple Lie superalgebras and their universal R-matrices, Publ. Res. Inst. Math. Sci. 30-1 (1994), 15-87.

[31] R.B. Zhang, Structure and representations of the quantum general linear supergroup, Comm. Math. Phys. 195-3 (1998), 525-547, online available arXiv:q-alg/9611024.

D.T.: Institut de recherche en mathématique et physique (IRMP), Université catholique de Louvain, Chemin du Cyclotron 2, building L7.01.02, 1348 Louvain-la-Neuve, Belgium
E-mail address: `daniel.tubbenhauer@uclouvain.be`

P.V.: Institut de recherche en mathématique et physique (IRMP), Université catholique de Louvain, Chemin du Cyclotron 2, building L7.01.02, 1348 Louvain-la-Neuve, Belgium
E-mail address: `pedro.vaz@uclouvain.be`

P.W.: Centre for Mathematical Sciences, University of Cambridge, Wilberforce Road, Cambridge CB3 0WB, United Kingdom
E-mail address: `p.wedrich@gmail.com`